

Graduate Aptitude Test in Engineering 2017

Question Paper Name: Mathematics 5th Feb 2017
Subject Name: Mathematics
Duration: 180
Total Marks: 100



Organizing Institute:

Indian Institute of Technology Roorkee



Question Number : 1**Correct : 1 Wrong : 0**

Consider the vector space $V = \{a_0 + a_1x + a_2x^2 : a_i \in \mathbb{R} \text{ for } i = 0, 1, 2\}$ of polynomials of degree at most 2. Let $f: V \rightarrow \mathbb{R}$ be a linear functional such that $f(1+x) = 0$, $f(1-x^2) = 0$ and $f(x^2-x) = 2$. Then $f(1+x+x^2)$ equals _____.

Question Number : 2**Correct : 1 Wrong : 0**

Let A be a 7×7 matrix such that $2A^2 - A^4 = I$, where I is the identity matrix. If A has two distinct eigenvalues and each eigenvalue has geometric multiplicity 3, then the total number of nonzero entries in the Jordan canonical form of A equals _____.

Question Number : 3**Correct : 1 Wrong : -0.33**

Let $f(z) = (x^2 + y^2) + i2xy$ and $g(z) = 2xy + i(y^2 - x^2)$ for $z = x + iy \in \mathbb{C}$. Then, in the complex plane $\mathbb{C}$,

- (A) f is analytic and g is NOT analytic
- (B) f is NOT analytic and g is analytic
- (C) neither f nor g is analytic
- (D) both f and g are analytic

Question Number : 4**Correct : 1 Wrong : 0**

If $\sum_{n=-\infty}^{\infty} a_n(z-2)^n$ is the Laurent series of the function $f(z) = \frac{z^4 + z^3 + z^2}{(z-2)^3}$ for $z \in \mathbb{C} \setminus \{2\}$, then a_{-2} equals _____.

Question Number : 5**Correct : 1 Wrong : -0.33**

Let $f_n: [0,1] \rightarrow \mathbb{R}$ be given by $f_n(x) = \frac{2x^2}{x^2 + (1-2nx)^2}$, $n = 1, 2, \dots$. Then the sequence (f_n)

- (A) converges uniformly on $[0,1]$
- (B) does NOT converge uniformly on $[0,1]$ but has a subsequence that converges uniformly on $[0,1]$
- (C) does NOT converge pointwise on $[0,1]$
- (D) converges pointwise on $[0,1]$ but does NOT have a subsequence that converges uniformly on $[0,1]$

Question Number : 6**Correct : 1 Wrong : 0**

Let $C: x^2 + y^2 = 9$ be the circle in $\mathbb{R}^2$ oriented positively. Then

$\frac{1}{\pi} \oint_C (3y - e^{\cos x^2}) dx + (7x + \sqrt{y^4 + 11}) dy$ equals _____.

Question Number : 7**Correct : 1 Wrong : -0.33**

Consider the following statements:

(P): There exists an unbounded subset of $\mathbb{R}$ whose Lebesgue measure is equal to 5.

(Q): If $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous and $g: \mathbb{R} \rightarrow \mathbb{R}$ is such that $f = g$ almost everywhere on $\mathbb{R}$, then g must be continuous almost everywhere on $\mathbb{R}$.

Which of the above statements hold TRUE?

- (A) Both P and Q
- (B) Only P
- (C) Only Q
- (D) Neither P nor Q

Question Number : 8**Correct : 1 Wrong : -0.33**

If $x^3 y^2$ is an integrating factor of $(6y^2 + a xy) dx + (6xy + b x^2) dy = 0$, where $a, b \in \mathbb{R}$, then

- (A) $3a - 5b = 0$
- (B) $2a - b = 0$
- (C) $3a + 5b = 0$
- (D) $2a + b = 0$

Question Number : 9**Correct : 1 Wrong : 0**

If $x(t)$ and $y(t)$ are the solutions of the system $\frac{dx}{dt} = y$ and $\frac{dy}{dt} = -x$ with the initial conditions $x(0) = 1$ and $y(0) = 1$, then $x(\pi/2) + y(\pi/2)$ equals _____.

Question Number : 10**Correct : 1 Wrong : -0.33**

If $y = 3e^{2x} + e^{-2x} - \alpha x$ is the solution of the initial value problem $\frac{d^2 y}{dx^2} + \beta y = 4\alpha x$, $y(0) = 4$ and $\frac{dy}{dx}(0) = 1$, where $\alpha, \beta \in \mathbb{R}$, then

- (A) $\alpha = 3$ and $\beta = 4$ (B) $\alpha = 1$ and $\beta = 2$
(C) $\alpha = 3$ and $\beta = -4$ (D) $\alpha = 1$ and $\beta = -2$

Question Number : 11**Correct : 1 Wrong : 0**

Let G be a non-abelian group of order 125. Then the total number of elements in $Z(G) = \{x \in G : gx = xg \text{ for all } g \in G\}$ equals _____.

Question Number : 12**Correct : 1 Wrong : 0**

Let F_1 and F_2 be subfields of a finite field F consisting of 2^9 and 2^6 elements, respectively. Then the total number of elements in $F_1 \cap F_2$ equals _____.

Question Number : 13**Correct : 1 Wrong : 0**

Consider the normed linear space $\mathbb{R}^2$ equipped with the norm given by $\|(x, y)\| = |x| + |y|$ and the subspace $X = \{(x, y) \in \mathbb{R}^2 : x = y\}$. Let f be the linear functional on X given by $f(x, y) = 3x$. If $g(x, y) = \alpha x + \beta y$, $\alpha, \beta \in \mathbb{R}$, is a Hahn-Banach extension of f on $\mathbb{R}^2$, then $\alpha - \beta$ equals _____.

Question Number : 14**Correct : 1 Wrong : -0.33**

For $n \in \mathbb{Z}$, define $c_n = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} e^{i(n-i)x} dx$, where $i^2 = -1$. Then $\sum_{n \in \mathbb{Z}} |c_n|^2$ equals

- (A) $\cosh(\pi)$ (B) $\sinh(\pi)$ (C) $\cosh(2\pi)$ (D) $\sinh(2\pi)$

Question Number : 15**Correct : 1 Wrong : 0**

If the fourth order divided difference of $f(x) = \alpha x^4 + 5x^3 + 3x + 2$, $\alpha \in \mathbb{R}$, at the points 0.1, 0.2, 0.3, 0.4, 0.5 is 5, then α equals _____.

Question Number : 16**Correct : 1 Wrong : 0**

If the quadrature rule $\int_0^2 f(x) dx \approx c_1 f(0) + 3 f(c_2)$, where $c_1, c_2 \in \mathbb{R}$, is exact for all polynomials of degree ≤ 1 , then $c_1 + 3c_2$ equals _____.

Question Number : 17**Correct : 1 Wrong : -0.33**

If $u(x, y) = 1 + x + y + f(xy)$, where $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a differentiable function, then u satisfies

(A) $x \frac{\partial u}{\partial x} - y \frac{\partial u}{\partial y} = x^2 - y^2$

(B) $x \frac{\partial u}{\partial x} - y \frac{\partial u}{\partial y} = 0$

(C) $x \frac{\partial u}{\partial x} - y \frac{\partial u}{\partial y} = x - y$

(D) $y \frac{\partial u}{\partial x} - x \frac{\partial u}{\partial y} = x - y$

Question Number : 18**Correct : 1 Wrong : -0.33**

The partial differential equation $x \frac{\partial^2 u}{\partial x^2} + (x - y) \frac{\partial^2 u}{\partial x \partial y} - y \frac{\partial^2 u}{\partial y^2} + \frac{1}{4} \left(\frac{\partial u}{\partial y} - \frac{\partial u}{\partial x} \right) = 0$ is

(A) hyperbolic along the line $x + y = 0$

(B) elliptic along the line $x - y = 0$

(C) elliptic along the line $x + y = 0$

(D) parabolic along the line $x + y = 0$

Question Number : 19**Correct : 1 Wrong : -0.33**

Let X and Y be topological spaces and let $f: X \rightarrow Y$ be a continuous surjective function. Which one of the following statements is TRUE?

(A) If X is separable, then Y is separable

(B) If X is first countable, then Y is first countable

(C) If X is Hausdorff, then Y is Hausdorff

(D) If X is regular, then Y is regular

Question Number : 20**Correct : 1 Wrong : -0.33**

Consider the topology $\mathcal{T} = \{U \subseteq \mathbb{Z} : \mathbb{Z} \setminus U \text{ is finite or } 0 \notin U\}$ on $\mathbb{Z}$. Then, the topological space $(\mathbb{Z}, \mathcal{T})$ is

(A) compact but NOT connected

(B) connected but NOT compact

(C) both compact and connected

(D) neither compact nor connected

Question Number : 21**Correct : 1 Wrong : -0.33**

Let $F(x)$ be the distribution function of a random variable X . Consider the functions:

$$G_1(x) = (F(x))^3, \quad x \in \mathbb{R},$$

$$G_2(x) = 1 - (1 - F(x))^5, \quad x \in \mathbb{R}.$$

Which of the above functions are distribution functions?

(A) Neither G_1 nor G_2

(B) Only G_1

(C) Only G_2

(D) Both G_1 and G_2

Question Number : 22**Correct : 1 Wrong : -0.33**

Let $X_1, X_2, \dots, X_n$ ($n \geq 2$) be independent and identically distributed random variables with finite variance σ^2 and let $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$. Then the covariance between $\bar{X}$ and $X_1 - \bar{X}$ is

(A) 0

(B) $-\sigma^2$

(C) $-\frac{\sigma^2}{n}$

(D) $\frac{\sigma^2}{n}$

Question Number : 23**Correct : 1 Wrong : 0**

Let $X_1, X_2, \dots, X_n$ ($n \geq 2$) be a random sample from a $N(\mu, \sigma^2)$ population, where $\sigma^2 = 144$. The smallest n such that the length of the shortest 95% confidence interval for μ will not exceed 10 is _____.

Question Number : 24**Correct : 1 Wrong : -0.33**

Consider the linear programming problem (LPP):

$$\text{Maximize } 4x_1 + 6x_2$$

$$\text{Subject to } x_1 + x_2 \leq 8,$$

$$2x_1 + 3x_2 \geq 18,$$

$$x_1 \geq 6, \quad x_2 \text{ is unrestricted in sign.}$$

Then the LPP has

- (A) no optimal solution
- (B) only one basic feasible solution and that is optimal
- (C) more than one basic feasible solution and a unique optimal solution
- (D) infinitely many optimal solutions

Question Number : 25**Correct : 1 Wrong : -0.33**

For a linear programming problem (LPP) and its dual, which one of the following is NOT TRUE?

- (A) The dual of the dual is primal
- (B) If the primal LPP has an unbounded objective function, then the dual LPP is infeasible
- (C) If the primal LPP is infeasible, then the dual LPP must have unbounded objective function
- (D) If the primal LPP has a finite optimal solution, then the dual LPP also has a finite optimal solution

Question Number : 26**Correct : 2 Wrong : 0**

If U and V are the null spaces of $\begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & 2 & 3 & 2 \\ 0 & 1 & 2 & 1 \end{bmatrix}$, respectively, then the dimension of the subspace $U + V$ equals _____.

Question Number : 27**Correct : 2 Wrong : -0.66**

Given two $n \times n$ matrices A and B with entries in $\mathbb{C}$, consider the following statements:

(P): If A and B have the same minimal polynomial, then A is similar to B .

(Q): If A has n distinct eigenvalues, then there exists $u \in \mathbb{C}^n$ such that $u, Au, \dots, A^{n-1}u$ are linearly independent.

Which of the above statements hold TRUE?

(A) Both P and Q

(B) Only P

(C) Only Q

(D) Neither P nor Q

Question Number : 28**Correct : 2 Wrong : 0**

Let $A = (a_{ij})$ be a 10×10 matrix such that $a_{ij} = 1$ for $i \neq j$ and $a_{ii} = \alpha + 1$, where $\alpha > 0$. Let λ and μ be the largest and the smallest eigenvalues of A , respectively. If $\lambda + \mu = 24$, then α equals _____.

Question Number : 29**Correct : 2 Wrong : 0**

Let C be the simple, positively oriented circle of radius 2 centered at the origin in the complex plane. Then

$$\frac{2}{\pi i} \int_C \left(z e^{(1/z)} + \tan\left(\frac{z}{2}\right) + \frac{1}{(z-1)(z-3)^2} \right) dz \text{ equals } \underline{\hspace{2cm}}.$$

Question Number : 30**Correct : 2 Wrong : -0.66**

Let $\operatorname{Re}(z)$ and $\operatorname{Im}(z)$, respectively, denote the real part and the imaginary part of a complex number z . Let $T: \mathbb{C} \cup \{\infty\} \rightarrow \mathbb{C} \cup \{\infty\}$ be the bilinear transformation such that $T(6) = 0$, $T(3 - 3i) = i$ and $T(0) = \infty$. Then, the image of $D = \{z \in \mathbb{C} : |z - 3| < 3\}$ under the mapping $w = T(z)$ is

- (A) $\{w \in \mathbb{C} : \operatorname{Im}(w) < 0\}$ (B) $\{w \in \mathbb{C} : \operatorname{Re}(w) < 0\}$
(C) $\{w \in \mathbb{C} : \operatorname{Im}(w) > 0\}$ (D) $\{w \in \mathbb{C} : \operatorname{Re}(w) > 0\}$

Question Number : 31**Correct : 2 Wrong : -0.66**

Let (x_n) and (y_n) be two sequences in a complete metric space (X, d) such that $d(x_n, x_{n+1}) \leq \frac{1}{n^2}$ and $d(y_n, y_{n+1}) \leq \frac{1}{n}$ for all $n \in \mathbb{N}$. Then

- (A) both (x_n) and (y_n) converge
(B) (x_n) converges but (y_n) need NOT converge
(C) (y_n) converges but (x_n) need NOT converge
(D) neither (x_n) nor (y_n) converges

Question Number : 32**Correct : 2 Wrong : 0**

Let $f: [0, 1] \rightarrow \mathbb{R}$ be given by $f(x) = 0$ if x is rational, and if x is irrational then $f(x) = 9^n$, where n is the number of zeroes immediately after the decimal point in the decimal representation of x . Then the Lebesgue integral $\int_0^1 f(x) dx$ equals _____.

Question Number : 33

Correct : 2 Wrong : -0.66

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $f(x, y) = \begin{cases} \sin\left(\frac{y^2}{x}\right) \sqrt{x^2 + y^2} & , \quad x \neq 0, \\ 0, & x = 0. \end{cases}$

Then, at $(0, 0)$,

- (A) f is continuous and the directional derivative of f does NOT exist in some direction
- (B) f is NOT continuous and the directional derivatives of f exist in all directions
- (C) f is NOT differentiable and the directional derivatives of f exist in all directions
- (D) f is differentiable

Question Number : 34

Correct : 2 Wrong : 0

Let D be the region in $\mathbb{R}^2$ bounded by the parabola $y^2 = 2x$ and the line $y = x$. Then $\iint_D 3xy \, dx \, dy$ equals _____.

Question Number : 35

Correct : 2 Wrong : -0.66

Let $y_1(x) = x^3$ and $y_2(x) = x^2|x|$ for $x \in \mathbb{R}$.

Consider the following statements:

- (P): $y_1(x)$ and $y_2(x)$ are linearly independent solutions of $x^2 \frac{d^2 y}{dx^2} - 4x \frac{dy}{dx} + 6y = 0$ on $\mathbb{R}$.
- (Q): The Wronskian $y_1(x) \frac{dy_2}{dx}(x) - y_2(x) \frac{dy_1}{dx}(x) = 0$ for all $x \in \mathbb{R}$.

Which of the above statements hold TRUE?

- (A) Both P and Q
- (B) Only P
- (C) Only Q
- (D) Neither P nor Q

Question Number : 36**Correct : 2 Wrong : 0**

Let α and β with $\alpha > \beta$ be the roots of the indicial equation of

$$(x^2 - 1)^2 \frac{d^2 y}{dx^2} + (x + 1) \frac{dy}{dx} - y = 0 \text{ at } x = -1. \text{ Then } \alpha - 4\beta \text{ equals } \underline{\hspace{2cm}}.$$

Question Number : 37**Correct : 2 Wrong : 0**

Let S_9 be the group of all permutations of the set $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$. Then the total number of elements of S_9 that commute with $\tau = (1\ 2\ 3)(4\ 5\ 6\ 7)$ in S_9 equals $\underline{\hspace{2cm}}$.

Question Number : 38**Correct : 2 Wrong : 0**

Let $\mathbb{Q}[x]$ be the ring of polynomials over $\mathbb{Q}$. Then the total number of maximal ideals in the quotient ring $\mathbb{Q}[x]/(x^4 - 1)$ equals $\underline{\hspace{2cm}}$.

Question Number : 39**Correct : 2 Wrong : -0.66**

Let $\{e_n : n \in \mathbb{N}\}$ be an orthonormal basis of a Hilbert space H . Let $T : H \rightarrow H$ be given by

$$Tx = \sum_{n=1}^{\infty} \frac{1}{n} \langle x, e_n \rangle e_n. \text{ For each } n \in \mathbb{N}, \text{ define } T_n : H \rightarrow H \text{ by } T_n x = \sum_{j=1}^n \frac{1}{j} \langle x, e_j \rangle e_j.$$

Then

- (A) $\|T_n - T\| \rightarrow 0$ as $n \rightarrow \infty$
- (B) $\|T_n - T\| \not\rightarrow 0$ as $n \rightarrow \infty$ but for each $x \in H$, $\|T_n x - Tx\| \rightarrow 0$ as $n \rightarrow \infty$
- (C) for each $x \in H$, $\|T_n x - Tx\| \rightarrow 0$ as $n \rightarrow \infty$ but the sequence $(\|T_n\|)$ is unbounded
- (D) there exist $x, y \in H$ such that $\langle T_n x, y \rangle \not\rightarrow \langle Tx, y \rangle$ as $n \rightarrow \infty$

Question Number : 40**Correct : 2 Wrong : -0.66**

Consider the subspace $V = \{(x_n) \in \ell^2 : \sum_{n=1}^{\infty} |x_n| < \infty\}$ of the Hilbert space ℓ^2 of all square summable real sequences. For $n \in \mathbb{N}$, define $T_n : V \rightarrow \mathbb{R}$ by $T_n((x_k)) = \sum_{i=1}^n x_i$.

Consider the following statements:

(P): $\{T_n : n \in \mathbb{N}\}$ is pointwise bounded on V .

(Q): $\{T_n : n \in \mathbb{N}\}$ is uniformly bounded on $\{x \in V : \|x\|_2 = 1\}$.

Which of the above statements hold TRUE?

(A) Both P and Q

(B) Only P

(C) Only Q

(D) Neither P nor Q

Question Number : 41**Correct : 2 Wrong : 0**

Let $p(x)$ be the polynomial of degree at most 2 that interpolates the data $(-1, 2)$, $(0, 1)$ and $(1, 2)$. If $q(x)$ is a polynomial of degree at most 3 such that $p(x) + q(x)$ interpolates the data $(-1, 2)$, $(0, 1)$, $(1, 2)$ and $(2, 11)$, then $q(3)$ equals _____.

Question Number : 42**Correct : 2 Wrong : -0.66**

Let J be the Jacobi iteration matrix of the linear system $\begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 2 \\ -4 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$.

Consider the following statements:

(P): One of the eigenvalues of J lies in the interval $[2, 3]$.

(Q): The Jacobi iteration converges for the above system.

Which of the above statements hold TRUE?

(A) Both P and Q

(B) Only P

(C) Only Q

(D) Neither P nor Q

Question Number : 43**Correct : 2 Wrong : 0**

Let $u(x, y)$ be the solution of $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 4u$ satisfying the condition $u(x, y) = 1$ on the circle $x^2 + y^2 = 1$. Then $u(2, 2)$ equals _____.

Question Number : 44**Correct : 2 Wrong : -0.66**

Let $u(r, \theta)$ be the bounded solution of the following boundary value problem in polar coordinates:

$$r^2 \frac{\partial^2 u}{\partial r^2} + r \frac{\partial u}{\partial r} + \frac{\partial^2 u}{\partial \theta^2} = 0, \quad 0 < r < 2 \quad \text{and} \quad 0 \leq \theta \leq 2\pi,$$

$$u(2, \theta) = \cos^2 \theta, \quad 0 \leq \theta \leq 2\pi.$$

Then $u(1, \pi/2) + u(1, \pi/4)$ equals

- (A) 1 (B) $\frac{9}{8}$ (C) $\frac{7}{8}$ (D) $\frac{3}{8}$

Question Number : 45**Correct : 2 Wrong : -0.66**

Let T_u and T_d denote the usual topology and the discrete topology on $\mathbb{R}$, respectively.

Consider the following three topologies:

T_1 = Usual topology on $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$,

T_2 = Topology generated by the basis $\{U \times V : U \in T_d, V \in T_u\}$ on $\mathbb{R} \times \mathbb{R}$,

T_3 = Dictionary order topology on $\mathbb{R} \times \mathbb{R}$.

Then

- (A) $T_3 \subsetneq T_1 \subseteq T_2$ (B) $T_1 \subsetneq T_2 \subsetneq T_3$
 (C) $T_3 \subseteq T_2 \subsetneq T_1$ (D) $T_1 \subsetneq T_2 = T_3$

Question Number : 46**Correct : 2 Wrong : 0**

Let X be a random variable with probability mass function

$$p(n) = \left(\frac{3}{4}\right)^{n-1} \left(\frac{1}{4}\right) \quad \text{for } n = 1, 2, \dots. \quad \text{Then } E(X - 3 | X > 3) \text{ equals } \underline{\hspace{2cm}}.$$

Question Number : 47**Correct : 2 Wrong : 0**

Let X and Y be independent and identically distributed random variables with probability mass function $p(n) = 2^{-n}$, $n = 1, 2, \dots$.

Then $P(X \geq 2Y)$ equals (rounded to 2 decimal places) _____.

Question Number : 48**Correct : 2 Wrong : 0**

Let $X_1, X_2, \dots$ be a sequence of independent and identically distributed Poisson random variables

with mean 4. Then $\lim_{n \rightarrow \infty} P\left(4 - \frac{2}{\sqrt{n}} < \frac{1}{n} \sum_{i=1}^n X_i < 4 + \frac{2}{\sqrt{n}}\right)$ equals _____.

Question Number : 49**Correct : 2 Wrong : 0**

Let X and Y be independent and identically distributed exponential random variables with probability density function

$$f(x) = \begin{cases} e^{-x}, & x > 0, \\ 0, & \text{otherwise.} \end{cases}$$

Then $P(\max(X, Y) < 2)$ equals (rounded to 2 decimal places) _____.

Question Number : 50**Correct : 2 Wrong : 0**

Let E and F be any two events with $P(E) = 0.4$, $P(F) = 0.3$ and $P(F | E) = 3 P(F | E^c)$.

Then $P(E | F)$ equals (rounded to 2 decimal places) _____.

Question Number : 51**Correct : 2 Wrong : -0.66**

Let $X_1, X_2, \dots, X_m$ ($m \geq 2$) be a random sample from a binomial distribution with parameters $n = 1$ and p , $p \in (0, 1)$, and let $\bar{X} = \frac{1}{m} \sum_{i=1}^m X_i$. Then a uniformly minimum variance unbiased estimator for $p(1-p)$ is

(A) $\frac{m}{m-1} \bar{X} (1 - \bar{X})$

(B) $\bar{X} (1 - \bar{X})$

(C) $\frac{m-1}{m} \bar{X} (1 - \bar{X})$

(D) $\frac{1}{m} \bar{X} (1 - m\bar{X})$

Question Number : 52**Correct : 2 Wrong : 0**

Let $X_1, X_2, \dots, X_9$ be a random sample from a $N(0, \sigma^2)$ population. For testing $H_0 : \sigma^2 = 2$ against $H_1 : \sigma^2 = 1$, the most powerful test rejects H_0 if $\sum_{i=1}^9 X_i^2 < c$, where c is to be chosen such that the level of significance is 0.1. Then the power of this test equals _____.

Question Number : 53**Correct : 2 Wrong : -0.66**

Let $X_1, X_2, \dots, X_n$ ($n \geq 2$) be a random sample from a $N(\theta, \theta)$ population, where $\theta > 0$, and let $W = \frac{1}{n} \sum_{i=1}^n X_i^2$. Then the maximum likelihood estimator of θ is

(A) $\frac{1}{2} + \frac{1}{2} \sqrt{1 - 4W}$

(B) $\frac{1}{2} + \frac{1}{2} \sqrt{1 + 4W}$

(C) $\frac{-1}{2} + \frac{1}{2} \sqrt{1 - 4W}$

(D) $\frac{-1}{2} + \frac{1}{2} \sqrt{1 + 4W}$

Question Number : 54

Correct : 2 Wrong : 0

Consider the following transportation problem. The entries inside the cells denote per unit cost of transportation from the origins to the destinations.

		Destination			Supply
		1	2	3	
Origin	1	4	3	6	20
	2	7	10	5	30
	3	8	9	7	50
Demand		10	30	60	

The optimal cost of transportation equals _____.

Question Number : 55

Correct : 2 Wrong : 0

Consider the linear programming problem (LPP):

$$\text{Maximize } kx_1 + 5x_2$$

$$\text{Subject to } x_1 + x_2 \leq 1,$$

$$2x_1 + 3x_2 \leq 1,$$

$$x_1, x_2 \geq 0.$$

If $x^* = (x_1^*, x_2^*)$ is an optimal solution of the above LPP with $k = 2$, then the largest value of k (rounded to 2 decimal places) for which x^* remains optimal equals _____.

Question Number : 56**Correct : 1 Wrong : -0.33**

The ninth and the tenth of this month are Monday and Tuesday _____.

- (A) figuratively (B) retrospectively (C) respectively (D) rightfully

Question Number : 57**Correct : 1 Wrong : -0.33**

It is _____ to read this year's textbook _____ the last year's.

- (A) easier, than (B) most easy, than (C) easier, from (D) easiest, from

Question Number : 58**Correct : 1 Wrong : -0.33**

A rule states that in order to drink beer, one must be over 18 years old. In a bar, there are 4 people. P is 16 years old, Q is 25 years old, R is drinking milkshake and S is drinking a beer. What must be checked to ensure that the rule is being followed?

- (A) Only P's drink
(B) Only P's drink and S's age
(C) Only S's age
(D) Only P's drink, Q's drink and S's age

Question Number : 59**Correct : 1 Wrong : -0.33**

Fatima starts from point P, goes North for 3 km, and then East for 4 km to reach point Q. She then turns to face point P and goes 15 km in that direction. She then goes North for 6 km. How far is she from point P, and in which direction should she go to reach point P?

- (A) 8 km, East (B) 12 km, North (C) 6 km, East (D) 10 km, North

Question Number : 60**Correct : 1 Wrong : -0.33**

500 students are taking one or more courses out of Chemistry, Physics, and Mathematics. Registration records indicate course enrolment as follows: Chemistry (329), Physics (186), Mathematics (295), Chemistry and Physics (83), Chemistry and Mathematics (217), and Physics and Mathematics (63). How many students are taking all 3 subjects?

- (A) 37 (B) 43 (C) 47 (D) 53

Question Number : 61**Correct : 2 Wrong : -0.66**

“If you are looking for a history of India, or for an account of the rise and fall of the British Raj, or for the reason of the cleaving of the subcontinent into two mutually antagonistic parts and the effects this mutilation will have in the respective sections, and ultimately on Asia, you will not find it in these pages; for though I have spent a lifetime in the country, I lived too near the seat of events, and was too intimately associated with the actors, to get the perspective needed for the impartial recording of these matters.”

Which of the following statements best reflects the author’s opinion?

- (A) An intimate association does not allow for the necessary perspective.
- (B) Matters are recorded with an impartial perspective.
- (C) An intimate association offers an impartial perspective.
- (D) Actors are typically associated with the impartial recording of matters.

Question Number : 62**Correct : 2 Wrong : -0.66**

Each of P, Q, R, S, W, X, Y and Z has been married at most once. X and Y are married and have two children P and Q. Z is the grandfather of the daughter S of P. Further, Z and W are married and are parents of R. Which one of the following must necessarily be FALSE?

- (A) X is the mother-in-law of R
- (B) P and R are not married to each other
- (C) P is a son of X and Y
- (D) Q cannot be married to R

Question Number : 63**Correct : 2 Wrong : -0.66**

1200 men and 500 women can build a bridge in 2 weeks. 900 men and 250 women will take 3 weeks to build the same bridge. How many men will be needed to build the bridge in one week?

- (A) 3000
- (B) 3300
- (C) 3600
- (D) 3900


Question Number : 64**Correct : 2 Wrong : -0.66**

The number of 3-digit numbers such that the digit 1 is never to the immediate right of 2 is

- (A) 781
- (B) 791
- (C) 881
- (D) 891

Question Number : 65 Correct : 2 Wrong : -0.66

A contour line joins locations having the same height above the mean sea level. The following is a contour plot of a geographical region. Contour lines are shown at 25 m intervals in this plot.



Which of the following is the steepest path leaving from P?

- (A) P to Q (B) P to R (C) P to S (D) P to T

Q. No.	Type	Section	Key	Marks
1	NAT	MA	1 to 1	1
2	NAT	MA	8 to 8	1
3	MCQ	MA	B	1
4	NAT	MA	48 to 48	1
5	MCQ	MA	D	1
6	NAT	MA	36 to 36	1
7	MCQ	MA	B	1
8	MCQ	MA	A	1
9	NAT	MA	0 to 0	1
10	MCQ	MA	C	1
11	NAT	MA	5 to 5	1
12	NAT	MA	8 to 8	1
13	NAT	MA	0 to 0	1
14	MCQ	MA	D	1
15	NAT	MA	5 to 5	1
16	NAT	MA	1 to 1	1
17	MCQ	MA	C	1
18	MCQ	MA	D	1
19	MCQ	MA	A	1
20	MCQ	MA	A	1
21	MCQ	MA	D	1
22	MCQ	MA	A	1
23	NAT	MA	23 to 23	1
24	MCQ	MA	B	1
25	MCQ	MA	C	1
26	NAT	MA	3 to 3	2
27	MCQ	MA	C	2
28	NAT	MA	7 to 7	2
29	NAT	MA	3 to 3	2
30	MCQ	MA	D	2
31	MCQ	MA	B	2
32	NAT	MA	9 to 9	2
33	MCQ	MA	C	2
34	NAT	MA	2 to 2	2
35	MCQ	MA	A	2
36	NAT	MA	2 to 2	2

37	NAT	MA	24 to 24	2
38	NAT	MA	3 to 3	2
39	MCQ	MA	A	2
40	MCQ	MA	B	2
41	NAT	MA	24 to 24	2
42	MCQ	MA	B	2
43	NAT	MA	64 to 64	2
44	MCQ	MA	C	2
45	MCQ	MA	D	2
46	NAT	MA	4 to 4	2
47	NAT	MA	0.27 to 0.30	2
48	NAT	MA	0.67 to 0.70	2
49	NAT	MA	0.73 to 0.77	2
50	NAT	MA	0.65 to 0.68	2
51	MCQ	MA	A	2
52	NAT	MA	0.49 to 0.51	2
53	MCQ	MA	D	2
54	NAT	MA	590 to 590	2
55	NAT	MA	3.32 to 3.34	2
56	MCQ	GA	C	1
57	MCQ	GA	A	1
58	MCQ	GA	B	1
59	MCQ	GA	A	1
60	MCQ	GA	D	1
61	MCQ	GA	A	2
62	MCQ	GA	D	2
63	MCQ	GA	C	2
64	MCQ	GA	C	2
65	MCQ	GA	B	2