

Andhra Pradesh State Council of Higher Education

Notations :

- 1.Options shown in green color and with ✓ icon are correct.
- 2.Options shown in red color and with ✗ icon are incorrect.

Question Paper Name :	BioTechnology 08th May 2024 Shift 1
Duration :	180
Total Marks :	200
Display Marks:	No
Share Answer Key With Delivery Engine :	Yes
Calculator :	None
Magnifying Glass Required? :	No
Ruler Required? :	No
Eraser Required? :	No
Scratch Pad Required? :	No
Rough Sketch/Notepad Required? :	No
Protractor Required? :	No
Show Watermark on Console? :	Yes
Highlighter :	No
Auto Save on Console?	Yes
Change Font Color :	No
Change Background Color :	No
Change Theme :	No
Help Button :	No
Show Reports :	No

Show Progress Bar :	No
Is this Group for Examiner? :	No
Examiner permission :	Cant View
Show Progress Bar? :	No

Mathematics

Section Id :	210688143
Section Number :	1
Mandatory or Optional :	Mandatory
Number of Questions :	50
Section Marks :	50
Enable Mark as Answered Mark for Review and Clear Response :	Yes
Maximum Instruction Time :	0
Is Section Default? :	null

Question Number : 1 Question Id : 2106887205 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If each element of a row or column of a determinant is multiplied by a constant K then the value of the determinant is

Options :

- 1. ✖ Added by k
- 2. ✔ Multiplied by k
- 3. ✖ Subtracted by k

4. ✖ Divided by k.

Question Number : 2 Question Id : 2106887206 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If $A = \begin{bmatrix} 1 & 2 & 3 \\ -2 & 1 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 3 & 1 \\ 5 & 4 & 2 \\ 1 & 5 & 3 \end{bmatrix}$ then $AB =$

Options :

1. ✖ $\begin{bmatrix} 15 & 26 & 4 \end{bmatrix}$

2. ✔ $\begin{bmatrix} 15 & 26 & 14 \\ 5 & 18 & 12 \end{bmatrix}$

3. ✖ $\begin{bmatrix} 15 & 5 \\ 26 & 18 \\ 14 & 12 \end{bmatrix}$

4. ✖ BA

Question Number : 3 Question Id : 2106887207 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The elements on the main diagonal of a skew symmetric matrix are all

Options :

1. ✓ zeros

2. ✗ One's

3. ✗ Unequal

4. ✗ >1

Question Number : 4 Question Id : 2106887208 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If ω is one of the imaginary cube roots of unity, find the value of the determinant

$$\begin{vmatrix} 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \end{vmatrix} =$$

Options :

1. ✓ zero

2. ✗ one

3. ✗ ω^2

4. ✗ ω

Question Number : 5 Question Id : 2106887209 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Every square matrix can be written as the sum of

Options :

1. ✖ Diagonal matrix & square matrix
2. ✖ Two rectangular matrices
3. ✖ Square and non-square matrices
4. ✔ Symmetric and skew symmetric matrix

Question Number : 6 Question Id : 2106887210 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

An improper fraction can be reduced to proper fraction by

Options :

1. ✖ Multiplication
2. ✔ Division

3. ✖ subtraction

4. ✖ Addition

Question Number : 7 Question Id : 2106887211 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\frac{x}{(x+2)(x-3)} =$$

Options :

1. ✖ $\frac{2}{5(x+2)} + \frac{3}{5(x-2)}$

2. ✖ $\frac{2}{5(x+2)} - \frac{3}{5(x-3)}$

3. ✔ $\frac{2}{5(x+2)} + \frac{3}{5(x-3)}$

4. ✖ $\frac{2}{5(x-3)} + \frac{3}{5(x+2)}$

Question Number : 8 Question Id : 2106887212 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The value of $\sin 210^\circ$

Options :

1. ✖ $\frac{1}{2}$

2. ✔ $-\frac{1}{2}$

3. ✖ $\frac{1}{\sqrt{2}}$

4. ✖ $-\frac{1}{\sqrt{2}}$

Question Number : 9 Question Id : 2106887213 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\cos n\pi =$$

Options :

1. ✖ -1

2. ✖ $-n$

3. ✔ $(-1)^n$

4. ✖ $(n)^{-1}$

Question Number : 10 Question Id : 2106887214 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$a \neq 0 \neq b, \sin x + \sin y = a, \cos x + \cos y = b$ then $\tan \frac{x+y}{2} =$

Options :

1. ✖ $\frac{b}{a}$

2. ✔ $\frac{a}{b}$

3. ✖ $\frac{a+b}{2}$

4. ✖ $\frac{a-b}{2}$

Question Number : 11 Question Id : 2106887215 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$f(x)$ is a periodic function of period k then the period of periodic function $f(ax+b)$ is

Options :

1. ✖ $\frac{k}{a}, a \neq 0$

2. ✖ $\frac{ak}{|b|}, b \neq 0$

3. ✖ $\frac{k+b}{a}, a \neq 0$

4. ✔ $\frac{k}{|a|}, a \neq 0$

Question Number : 12 Question Id : 2106887216 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If $7\sin^2\theta + 3\cos^2\theta = 4$, then $\theta =$

Options :

1. ✖ $\pm \frac{\pi}{3}$

2. ✔ $\pm \frac{\pi}{6}$

3. ✖ $\pm \frac{\pi}{4}$

4. ✖ $\pm \frac{\pi}{2}$

Question Number : 13 Question Id : 2106887217 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The range of $\cos^{-1}x$ is

Options :

1. ✓ $[0, \pi]$

2. ✗ $[-\pi, \pi]$

3. ✗ $[0, -\pi]$

4. ✗ $(0, \pi)$

Question Number : 14 Question Id : 2106887218 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Assume $x > 0, y > 0$. Then which one of the following is true ?

Options :

1. ✓ If $xy < 1$ then $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right)$

2. ✗ If $xy > 1$ then $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right)$

3. ✖ If $xy=1$ then $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right)$

4. ✖ If $xy=1$ then $\tan^{-1}x - \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right)$

Question Number : 15 Question Id : 2106887219 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In $\triangle ABC$ $(a+b+c)(b+c-a)=3bc$, then angle A =

Options :

1. ✖ 90°

2. ✖ 120°

3. ✔ 60°

4. ✖ 45°

Question Number : 16 Question Id : 2106887220 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In $\triangle ABC$, $\tan \frac{A}{2} = \frac{5}{6}$, $\tan \frac{C}{2} = \frac{2}{5}$ then a,b,c are in

Options :

1. ✖ Geometric progression
2. ✔ Arithmetic progression
3. ✖ Harmonic progression
4. ✖ Arithmetico – Geometric progression

Question Number : 17 Question Id : 2106887221 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In any ΔABC , $\tan \frac{B-C}{2} =$

Options :

1. ✖ $b \pm c \cot \frac{A}{2}$

2. ✔ $\frac{b-c}{b+c} \cot \frac{A}{2}$

3. ✖ $(b-c) \tan \frac{A}{2}$

4. ✖ $\tan \frac{C}{2}$

Question Number : 18 Question Id : 2106887222 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Conjugate of $\frac{1-i}{1+i}$ is

Options :

1. ✖ $-3i$

2. ✖ $-i$

3. ✔ i

4. ✖ $6i$

Question Number : 19 Question Id : 2106887223 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Standard form of $(-1 + 2i) + \left(\frac{1}{2} - i\right)$ is

Options :

1. ✖ $\frac{1}{2} - i$

2. ✔ $-\frac{1}{2} + i$

3. ✖ $-\frac{1}{2} - i$

4. ✖ $\frac{1}{2} \pm i$

Question Number : 20 Question Id : 2106887224 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If the circle is $x^2 + y^2 + 6x - 8y + c = 0$ has radius 6 units, Then value of c is

Options :

1. ✔ -11

2. ✖ 11

3. ✖ 25

4. ✖ 6

Question Number : 21 Question Id : 2106887225 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The equation of the parabola whose focus is (8,0) and the vertex is (0,0) is

Options :

1. ✖ $y^2 = 12x$

2. ✖ $y^2 = x$

3. ✔ $y^2 = 32x$

4. ✖ $y^2 = 16x$

Question Number : 22 Question Id : 2106887226 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The eccentricity of the ellipse $x^2 + 2y^2 = 3$ is

Options :

1. ✖ $e = \frac{3}{\sqrt{2}}$

2. ✖ $e = \frac{1}{\sqrt{3}}$

3. ✖ $e = -\frac{1}{\sqrt{2}}$

4. ✔ $e = \frac{1}{\sqrt{2}}$

Question Number : 23 Question Id : 2106887227 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In the Ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$ the length of the latus rectum is _____

Options :

1. ✖ $\frac{2a^2}{b}$

2. ✔ $\frac{2b^2}{a}$

3. ✖ $\frac{2a^2}{b^2}$

4. ✖ $2ab$

Question Number : 24 Question Id : 2106887228 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The equation of the Hyperbola with foci $(\pm 2, 0)$ and eccentricity $3/2$ is

Options :

1. ✖ $\frac{9x^2}{16^2} + \frac{9y^2}{10^2} = 1$

2. ✔

$$\frac{x^2}{16/9} - \frac{y^2}{20/9} = 1$$

3. ✖ $\frac{x^2}{16^2} - \frac{y^2}{20^2} = 1$

4. ✖ $\frac{x^2}{2^2} - \frac{y^2}{20^2} = 1$

Question Number : 25 Question Id : 2106887229 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If the coordinates at one end of a diameter of the circle $x^2 + y^2 - 8x - 4y + c = 0$ are $(-3, 2)$ then the coordinates at the other end are

Options :

1. ✖ $(5, 11)$

2. ✖ $(6, 2)$

3. ✖ $(2, 11)$

4. ✔ $(11, 2)$

Question Number : 26 Question Id : 2106887230 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time : 0

If $a > 0$, then $\lim_{x \rightarrow 0} \frac{a^x - 1}{x} =$

Options :

1. ✖ $\log x$

2. ✖ 1

3. ✔ $\log a$

4. ✖ $\log\left(\frac{a}{x}\right)$

Question Number : 27 Question Id : 2106887231 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Differentiation of $\sin x^n$ with respect to x .

Options :

1. ✔ $nx^{n-1} \cos x^n$

2. ✖ $x^{n-1} \cos x^n$

3. ✖ $\cos x^n$

4. ✖

$$n \cos x^n$$

Question Number : 28 Question Id : 2106887232 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\frac{d}{dx} \left(\sin^{-1} \frac{x}{a} \right) =$$

Options :

1. ✓ $\frac{1}{\sqrt{a^2 - x^2}}$

2. ✗ $\frac{1}{\sqrt{a^2 + x^2}}$

3. ✗ $\frac{1}{\sqrt{x^2 - a^2}}$

4. ✗ $\frac{-1}{\sqrt{a^2 - x^2}}$

Question Number : 29 Question Id : 2106887233 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\frac{d}{dx} (e^{3 \log x}) =$$

Options :

1. ✖ $3x$

2. ✖ $3\log x$

3. ✖ $\log 3$

4. ✔ $3x^2$

Question Number : 30 Question Id : 2106887234 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\frac{d}{dx}[\log|x|] =$$

Options :

1. ✖ $\frac{1}{|x|}$

2. ✔ $\frac{1}{x}$

3. ✖ $|x|$

4. ✖ x

Question Number : 31 Question Id : 2106887235 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$y = \cos x$ then $\frac{d^2y}{dx^2}$ is

Options :

1. ✖ $\cos x$

2. ✖ $\sin x$

3. ✔ $-\cos x$

4. ✖ $-\sin x$

Question Number : 32 Question Id : 2106887236 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The angle between the curves $x^2 + 4y = 0, xy = 2$ is

Options :

1. ✔ $\tan^{-1} 3$

2. ✖ $\cot^{-1} 1$

3. ✖ $\tan^{-1} 4$

4. ✖ $\cot^{-1} 3$

Question Number : 33 Question Id : 2106887237 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The slope of the tangent to the curve $y = \frac{x-1}{x+1}$ at (0,1)

Options :

1. ✖ 4

2. ✖ -2

3. ✖ 5

4. ✔ 2

Question Number : 34 Question Id : 2106887238 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If $z = x^2 + y^2$ then $x \frac{\partial z}{\partial y} - y \frac{\partial z}{\partial x} =$

Options :

1. ✖ $2y-2x$

2. ✖ $2x+2y$

3. ✔ 0

4. ✖ $4xy$

Question Number : 35 Question Id : 2106887239 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$z = \frac{x^3+y^3}{x+y}$, is a homogeneous function of degree _____

Options :

1. ✔ 2

2. ✖ 3

3. ✖ 0

4. ✖ 1

Question Number : 36 Question Id : 2106887240 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\int (x^{2/3} + 1) dx =$$

Options :

1. ✓ $\frac{3}{5}x^{5/3} + x + c$

2. ✗ $\frac{5}{3}x^{5/3} + x + c$

3. ✗ $\frac{3}{5}x^{5/3} + c$

4. ✗ $\frac{3}{5}x^{3/5} + x + c$

Question Number : 37 Question Id : 2106887241 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\int \frac{dx}{x^2-16} =$$

Options :

1. ✗ $\frac{1}{16} \log \left| \frac{x-8}{x+4} \right| + c$

2. ✗ $\frac{1}{4} \log \left| \frac{x-4}{x+4} \right| + c$

3. ✓ $\frac{1}{8} \log \left| \frac{x-4}{x+4} \right| + c$

4. ✗ $\frac{1}{16} \log \left| \frac{x-4}{x+4} \right| + c$

Question Number : 38 Question Id : 2106887242 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\int \frac{\sin(\tan^{-1}x)dx}{1+x^2} =$$

Options :

1. ✗ $-\cos x + c$

2. ✓ $-\cos (\tan^{-1}x) + c$

3. ✗ $-\sin (\tan^{-1}x) + c$

4. ✗ $(\tan^{-1}x) + c$

Question Number : 39 Question Id : 2106887243 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\int \cos \frac{x}{2} dx =$$

Options :

1. ✖ $2 \cos \frac{x}{2} + c$

2. ✔ $2 \sin \frac{x}{2} + c$

3. ✖ $2 \sin 2x + c$

4. ✖ $-2 \sin \frac{x}{2} + c$

Question Number : 40 Question Id : 2106887244 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

$$\int e^x \cos x dx =$$

Options :

1. ✔ $\frac{1}{2} e^x (\cos x + \sin x) + c$

2. ✖ $\frac{1}{2} e^x (\cos x - \sin x) + c$

3. ✖ $\frac{1}{2}e^x \sin x + c$

4. ✖ $\frac{1}{2}(\cos x + \sin x) + c$

Question Number : 41 Question Id : 2106887245 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The area of the region bounded by the curve $y = f(x)$, x - axis and the lines $x = a$ and $x = b$ ($b > a$) is given by

Options :

1. ✖ $\int_b^a y dx$

2. ✖ $-\int_a^b y dx$

3. ✖ $\int_a^b x dy$

4. ✔ $\int_a^b y dx$

Question Number : 42 Question Id : 2106887246 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If $f(x)$ is an even function, then $\int_{-a}^a f(x)dx =$

Options :

1. ✖ $-\int_{-a}^a f(x)dx$

2. ✖ $2\int_{-a}^a f(x)dx$

3. ✔ $2\int_0^a f(x)dx$

4. ✖ $\int_0^a f(x)dx$

Question Number : 43 Question Id : 2106887247 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Find maxima (or) minima for the curve $y = 2x^4 - x^2$

Options :

1. ✔ 'y' is minimum at $x = \pm \frac{1}{2}$

2. ✖ 'y' is maximum for $x = -\frac{1}{4}$

3. ✖ 'y' is maximum for $x = \pm \frac{1}{2}$

4. ✖ 'y' is maximum for $x = +\frac{1}{4}$

Question Number : 44 Question Id : 2106887248 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Order of the differential equation $\left[\frac{d^2y}{dx^2} + \left(\frac{dy}{dx} \right)^3 \right]^{6/5} = 6y$ is

Options :

1. ✖ 3

2. ✔ 2

3. ✖ 5

4. ✖ 1

Question Number : 45 Question Id : 2106887249 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The general solution of the differential equation $\frac{dy}{dx} = \frac{1+y^2}{1+x^2}$ is

Options :

1. ✓ $\tan^{-1}y - \tan^{-1}x = c$

2. ✗ $\tan^{-1}y + \tan^{-1}x = c$

3. ✗ $\tan^{-1}y = c$

4. ✗ $\tan^{-1}y/x = c$

Question Number : 46 Question Id : 2106887250 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The differential equation representing the family of curves $y = mx$ where, m is arbitrary Constant is

Options :

1. ✗ $\frac{dy}{dx} - y = 0$

2. ✗ $\frac{dy}{dx} + y = 0$

3. ✓ $x \frac{dy}{dx} - y = 0$

4. ✖ $x dx - y dy = y$

Question Number : 47 Question Id : 2106887251 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which one of the statement is true?

Options :

1. ✖ Order of differential equation is the order of the lowest order derivative occurring in the differential equation.

2. ✖ A function which satisfies the given differential equation is not its solution .

3. ✖ An equation involving derivatives of the dependent variable with respect to dependent variable is known as a differential equation.

4. ✔ Degree of a differential equation is defined if it is a polynomial equation in its Derivatives.

Question Number : 48 Question Id : 2106887252 Display Question Number : Yes Is Question

Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The Integrating factor of the differential equation $x \frac{dy}{dx} + 2y = x^2 (x \neq 0)$ is

Options :

1. ✖ x

2. ✖ $\log x$

3. ✖ $x \log x$

4. ✔ x^2

Question Number : 49 Question Id : 2106887253 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The linear form of $x \log x \frac{dy}{dx} + y = 2 \log x$ is

Options :

1. ✖ $\frac{dy}{dx} - \frac{y}{x \log x} = \frac{1}{x}$

2. ✔ $\frac{dy}{dx} + \frac{y}{x \log x} = \frac{2}{x}$

3. ✖ $\frac{dy}{dx} + \frac{y}{x \log x} = \frac{1}{x}$

4. ✖ $\frac{dy}{dx} + \frac{y}{x \log x} = 1$

Question Number : 50 Question Id : 2106887254 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The particular integral of $\frac{d^2y}{dx^2} - 4y = e^{2x}$ is

Options :

1. ✖ $\frac{1}{4} e^{2x}$

2. ✖ $\frac{1}{4x} e^{2x}$

3. ✔ $\frac{1}{4} x e^{2x}$

4. ✖ 0

Physics

Section Id :

210688144

Section Number :	2
Mandatory or Optional :	Mandatory
Number of Questions :	25
Section Marks :	25
Enable Mark as Answered Mark for Review and Clear Response :	Yes
Maximum Instruction Time :	0
Is Section Default? :	null

Question Number : 51 Question Id : 2106887255 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

N Kg^{-1} is the unit of

Options :

1. ✖ Velocity
2. ✔ Acceleration
3. ✖ Force
4. ✖ Momentum

Question Number : 52 Question Id : 2106887256 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A system has basic dimensions as density 'D', velocity 'V' and area 'A'. The dimensional representation of force in this system is

Options :

1. ✓ $A V^2 D$

2. ✗ $A V D^2$

3. ✗ $A^2 V D$

4. ✗ $A^0 V^2 D$

Question Number : 53 Question Id : 2106887257 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If The magnitude of vectors **A**, **B** and **C** are 5, 4 and 3 units respectively and $\mathbf{A} = \mathbf{B} + \mathbf{C}$, then the angle between vectors **A** and **C** is

Options :

1. ✗ $\cos^{-1}(4/5)$

2. ✗ π

3. ✓ $\cos^{-1}(3/5)$

4. ✗ $\sin^{-1}(3/4)$

Question Number : 54 Question Id : 2106887258 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If the sum of two unit vectors is also a unit vector, then the magnitude of their difference is

Options :

1. ✖ 1

2. ✖ $\frac{1}{2}$

3. ✖ $\frac{1}{\sqrt{2}}$

4. ✔ $\sqrt{3}$

Question Number : 55 Question Id : 2106887259 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A particle starting from rest moves in a straight line with uniform acceleration a . The average velocity of the particle in first 's' distance is

Options :

1. ✔ $\sqrt{\frac{as}{2}}$

2. ✖ $\sqrt{\frac{3as}{2}}$

3. ✖ $\sqrt{2as}$

4. ✖ *as*

Question Number : 56 Question Id : 2106887260 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A projectile is thrown with speed u making angle θ with the horizontal at $t = 0$. It just crosses two points of equal height at time $t = 1\text{ s}$ and $t = 3\text{ s}$ respectively. The maximum height attained by the projectile is (take $g = 10\text{ ms}^{-2}$)

Options :

1. ✖ 10m

2. ✔ 20m

3. ✖ 15m

4. ✖ 22m

Question Number : 57 Question Id : 2106887261 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A body is falling from height 'H' takes time 'T' seconds to reach the ground. The time taken to cover the first half of height is

Options :

1. ✔

$\frac{T}{\sqrt{2}}$

2. ✖ $\sqrt{2} T$

3. ✖ $\sqrt{3} T$

4. ✖ $\frac{T}{\sqrt{3}}$

Question Number : 58 Question Id : 2106887262 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A body sliding on ice with a velocity 8 ms^{-1} comes to rest after travelling 40 m. The coefficient of friction between the body and ice is ($g = 10 \text{ ms}^{-2}$)

Options :

1. ✖ 0.02

2. ✖ 0.05

3. ✔ 0.08

4. ✖ 0.2

Question Number : 59 Question Id : 2106887263 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If a body placed on a rough inclined plane of gradient 1 in 4, just begins to slide, then coefficient of friction between the plane and body is

Options :

1. ✖ $\frac{2}{\sqrt{15}}$

2. ✖ $\frac{1}{\sqrt{2}}$

3. ✖ $\frac{1}{\sqrt{5}}$

4. ✔ $\frac{1}{\sqrt{15}}$

Question Number : 60 Question Id : 2106887264 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A cube of 10 N weight rests on a rough inclined plane of slope 3 in 5. If the coefficient of friction between plane and cube is 0.6, then minimum force required to start the cube moving up the plane is

Options :

1. ✖ 2N

2.

✖ 6N

3. ✔ 10.8N

4. ✖ 4.5N

Question Number : 61 Question Id : 2106887265 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A pump can take out 7200 Kg of water per hour from a 100 m deep well. If the efficiency of the pump is 50% then power of the pump is ($g = 10 \text{ ms}^{-2}$)

Options :

1. ✖ 2 KW

2. ✔ 4 KW

3. ✖ 7.2 KW

4. ✖ 3.6 KW

Question Number : 62 Question Id : 2106887266 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

When a force $\mathbf{F} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ acts on a body to move it from $\mathbf{r}_1 = \mathbf{i} + \mathbf{j} + \mathbf{k}$ to $\mathbf{r}_2 = \mathbf{i} - \mathbf{j} + 2\mathbf{k}$, then the work done by the force is

Options :

1. ✖ -3 J

2. ✔ -1 J

3. ✖ 2 J

4. ✖ 3 J

Question Number : 63 Question Id : 2106887267 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The K.E. of a body moving with a speed of 10 m/s is 30 J. If its speed becomes 30 m/s, then its K.E. will be

Options :

1. ✖ 10 J

2. ✖ 90 J

3. ✖ 180 J

4. ✔ 270 J

Question Number : 64 Question Id : 2106887268 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The maximum speed of a particle executing SHM is 1 m/s and maximum acceleration is 1.57 m/s^2 . Its time period is

Options :

1. ✓ 4 sec

2. ✗ 1.57 sec

3. ✗ 2 sec

4. ✗ $\frac{1}{1.57}$

Question Number : 65 Question Id : 2106887269 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A girl is swinging on a swing in the sitting position. If the girl stands up, the time period of the string will

Options :

1. ✗ Increase

2. ✓

Decrease

- 3. ✖ Remains same
- 4. ✖ Becomes erratic

Question Number : 66 Question Id : 2106887270 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A light spring supports 200 gm weight at its lower end; it oscillates with a period of 1 sec.
How much weight must be removed from the lower end to reduce the period to 0.5 sec?

Options :

- 1. ✖ 100 gm.
- 2. ✖ 50 gm.
- 3. ✔ 150 gm.
- 4. ✖ 200 gm.

Question Number : 67 Question Id : 2106887271 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The velocity of sound in any medium depends upon

Options :

1. ✖ Intensity and elasticity
2. ✖ Amplitude and density
3. ✔ elasticity and density
4. ✖ Amplitude and elasticity

Question Number : 68 Question Id : 2106887272 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The beat frequency produced by the vibrations of $x_1 = A \sin (320\pi t)$ and $x_2 = A \sin (326\pi t)$ is

Options :

1. ✖ 6
2. ✖ 4
3. ✖ 2
4. ✔ 3

Question Number : 69 Question Id : 2106887273 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The Boyle's law is stated by $PV = C$, C depends on

Options :

1. ✖ Nature of gas
2. ✖ Atomic weight of gas
3. ✖ Temperature of gas
4. ✔ Quantity and temperature of gas

Question Number : 70 Question Id : 2106887274 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The equation of state for 5g of oxygen(O_2) at pressure P and temperature T, when occupying a volume V, will be (R is universal gas constant)

Options :

1. ✖ $PV = 5RT$
2. ✖ $PV = \frac{5}{2} RT$
3. ✖

$$PV = \frac{5}{16} RT$$

4. ✓ $PV = \frac{5}{32} RT$

Question Number : 71 Question Id : 2106887275 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The volume of a gas at constant pressure of 10^3 N/m^2 expands by 0.25 m^3 . The work done in this process is

Options :

1. ✗ 25J

2. ✗ 50J

3. ✓ 250J

4. ✗ 5J

Question Number : 72 Question Id : 2106887276 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

For an adiabatic expansion of a perfect gas the value of $\frac{\Delta P}{P}$ is equal to

Options :

1. ✗

$$\frac{\Delta V}{V}$$

2. ✖ $\gamma \frac{\Delta V}{V}$

3. ✔ $-\gamma \frac{\Delta V}{V}$

4. ✖ $\gamma - \frac{\Delta V}{V}$

Question Number : 73 Question Id : 2106887277 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

First law of Thermodynamics is a special case of

Options :

1. ✖ Boyle's law

2. ✖ Charles law

3. ✖ Law of conservation of mass

4. ✔ Law of conservation of energy

Question Number : 74 Question Id : 2106887278 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If the critical angle for total internal reflection from a medium to vacuum is 30° , the velocity of light in the medium is

Options :

1. ✖ $3 \times 10^8 \text{ m/s}$
2. ✔ $1.5 \times 10^8 \text{ m/s}$
3. ✖ $\sqrt{3} \times 10^8 \text{ m/s}$
4. ✖ $2 \times 10^8 \text{ m/s}$

Question Number : 75 Question Id : 2106887279 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Light rays of wave length $4.36 \times 10^{-7} \text{ m}$ incident on a metal surface of work function 1.24 eV. The stopping potential required to stop the emission of photoelectrons is

Options :

1. ✔ 1.6 eV
2. ✖ 1.24 eV

3. ✖ 3.2 eV

4. ✖ 4.8 eV

Chemistry

Section Id :	210688145
Section Number :	3
Mandatory or Optional :	Mandatory
Number of Questions :	25
Section Marks :	25
Enable Mark as Answered Mark for Review and Clear Response :	Yes
Maximum Instruction Time :	0
Is Section Default? :	null

Question Number : 76 Question Id : 2106887280 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

According to Bohr's theory of hydrogen atom, the angular momentum of electron in fourth orbit of H-atom is equal to

Options :

1. ✖ $\frac{h}{2\pi}$

2. ✓ $\frac{2h}{\pi}$

3. ✗ $\frac{3h}{2\pi}$

4. ✗ $\frac{4h}{\pi}$

Question Number : 77 Question Id : 2106887281 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The quantum number which describes the shape of an atomic orbital is

Options :

1. ✓ Azimuthal Quantum Number

2. ✗ Principal Quantum Number

3. ✗ Spin Quantum Number

4. ✗ Magnetic Quantum Number

Question Number : 78 Question Id : 2106887282 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Identify the element in which the ratio of s-electrons to p-electrons is 3:5

Options :

1. ✖ P

2. ✖ Al

3. ✔ S

4. ✖ K

Question Number : 79 Question Id : 2106887283 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The pair of molecules in which the central atom has octet of electrons is

Options :

1. ✖ $\text{BeCl}_2, \text{BF}_3$

2. ✖ $\text{H}_2\text{O}, \text{BeCl}_2$

3. ✓ $\text{H}_2\text{O}, \text{NH}_3$

4. ✗ NH_3, BF_3

Question Number : 80 Question Id : 2106887284 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The electronic configuration of an element M is $[\text{Ne}]3\text{S}^1$ and that of element X is $[\text{He}]2\text{S}^22\text{P}^5$. The type of bond present between M and X is

Options :

1. ✗ Covalent Bond

2. ✓ Electrovalent Bond

3. ✗ Co-ordinate Covalent Bond

4. ✗ Hydrogen Bond

Question Number : 81 Question Id : 2106887285 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The absolute weight of one molecule of water (in g) is ($N_A=6 \times 10^{23} \text{ mol}^{-1}$)

Options :

1. ✖ 1.5×10^{-23}

2. ✔ 3.0×10^{-23}

3. ✖ 4.5×10^{-23}

4. ✖ 2.0×10^{-23}

Question Number : 82 Question Id : 2106887286 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The weight of sodium sulphate (molar mass 142 g mol^{-1}) required to prepare 500 ml of 0.03 M solution is

Options :

1. ✔ 2.13 g

2. ✖ 4.26 g

3. ✖ 1.065 g

4. ✖ 3.195 g

Question Number : 83 Question Id : 2106887287 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The number of H^+ ions present in 100 ml of 0.05 M H_2SO_4 solution is ($N_A = 6 \times 10^{23} \text{ mol}^{-1}$)

Options :

1. ✖ 6.0×10^{24}

2. ✖ 6.0×10^{22}

3. ✔ 6.0×10^{21}



4. ✖ 3.0×10^{23}

Question Number : 84 Question Id : 2106887288 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0


The conjugate acid and conjugate base of HCO_3^- are respectively

Options :


1. ✖



2. ✖



3. ✔



4. ✖

Question Number : 85 Question Id : 2106887289 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The pH of 0.005 M H_2SO_4 solution will be;

Options :

5

1. ✖

2

2. ✔

3

3. ✖

4

4. ✖

Question Number : 86 Question Id : 2106887290 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In an electrochemical cell, the electrons flow from

Options :

Cathode to Anode

1. ✖

Anode to Cathode

2. ✔

Anode to Solution

3. ✖

Solution to Cathode

4. ✖

Question Number : 87 Question Id : 2106887291 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

How many faradays are required to reduce 1 mole of MnO_4^- ions to Mn^{2+} ions?

Options :

1. ✔ 5

2. ✖

2

3. ✖ 4

4. ✖ 3

Question Number : 88 Question Id : 2106887292 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

At 298 K, the emf of the cell, $M|M^{2+}(1M) \parallel Cu^{2+}(1M)|Cu$ is 'x' V. If $E_{Cu^{2+}|Cu}^0 = +0.34V$,

then $E_{M^{2+}|M}^0$ (in V) is

Options :

1. ✖ $(x - 0.34)$ 2. ✔ $(0.34 - x)$ 3. ✖ $(0.34 + x)$ 4. ✖ $\frac{0.34}{x}$

Question Number : 89 Question Id : 2106887293 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Identify the strongest reducing agent from the following:

Options :

1. ✓ $E_{K^+|K}^0 = -2.93 \text{ V}$

2. ✗ $E_{Al^{3+}|Al}^0 = -1.66 \text{ V}$

3. ✗ $E_{Zn^{2+}|Zn}^0 = -0.76 \text{ V}$

4. ✗ $E_{Ag^+|Ag}^0 = +0.34 \text{ V}$

Question Number : 90 Question Id : 2106887294 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The formula of Zeolite can be represented as Na_2Z . The metal atom present in Z is

Options :



3. ✖ Mg

4. ✔ Al

Question Number : 91 Question Id : 2106887295 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following salts causes maximum hardness to water sample, when they are in equal amounts?

Options :

1. ✖ MgSO_4 (Molecular Weight = 120u)

2. ✔ MgCl_2 (Molecular Weight = 95u)

3. ✖ CaCl_2 (Molecular Weight = 111u)

4. ✖ $\text{Ca}(\text{HCO}_3)_2$ (Molecular Weight = 162u)

Question Number : 92 Question Id : 2106887296 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Permanent hardness of water cannot be removed by

Options :

1. ✓ Boiling the hard water
2. ✗ Treatment with washing soda
3. ✗ Passing through Zeolite
4. ✗ Passing through ion exchange resins

Question Number : 93 Question Id : 2106887297 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following statements is not correct about stress cells?

Options :

1. ✗ They are formed between different parts of the same metal
2. ✓ Stressed part of the metal acts as cathode
3. ✗ Stressed part of the metal acts as anode

4. ✖ Anodic part undergoes corrosion

Question Number : 94 Question Id : 2106887298 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Tarnishing of silver is due to the formation of

Options :

1. ✖ AgCl

2. ✖ Ag_2CO_3

3. ✖ Ag_2O

4. ✔ Ag_2S

Question Number : 95 Question Id : 2106887299 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following is not a natural polymer?

Options :

1. ✖ Wool

2. ✖ Cellulose

3. ✖ Strach

4. ✔ Rayon

Question Number : 96 Question Id : 2106887300 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Neoprene is an example of

Options :

1. ✔ Elastomer

2. ✖ Thermoplastic Polymer

3. ✖ Thermosetting Polymer

4. ✖ Co-Polymer

Question Number : 97 Question Id : 2106887301 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time : 0

The element that is added to raw rubber vulcanization is

Options :

1. ✓ S

2. ✗ Se

3. ✗ C

4. ✗ B

Question Number : 98 Question Id : 2106887302 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0


The major components of water gas are

Options :

1. ✓ H_2 , CO

2. ✗ H_2 , CO_2

3. ✗ CO, N_2



4. ✖

Question Number : 99 Question Id : 2106887303 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following is not a greenhouse gas?

Options :

1. ✖ O_3

2. ✖ CO_2

3. ✖ CH_4

4. ✔ N_2

Question Number : 100 Question Id : 2106887304 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The acid that is believed to be mainly responsible for the damage of Taj mahal is

Options :

1. ✔ H_2SO_4

2. ✖ HF

3. ✖ H₃PO₄

4. ✖ HCl

BioTechnology

Section Id :	210688146
Section Number :	4
Mandatory or Optional :	Mandatory
Number of Questions :	100
Section Marks :	100
Enable Mark as Answered Mark for Review and Clear Response :	Yes
Maximum Instruction Time :	0
Is Section Default? :	null

Question Number : 101 Question Id : 2106887305 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The replica plate technique is used for the isolation of

Options :

1. ✖ Auxotrophs

2. ✓ Revertants

3. ✗ Analog-resistant mutants

4. ✗ Prototrophs

Question Number : 102 Question Id : 2106887306 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following organism strain is improved by using parasexual cycle ?

Options :

1. ✓ *Aspergillus niger*

2. ✗ *Pencillium notatum*

3. ✗ *Brevibacterium flavum*

4. ✗ *Saccharomyces cerevisiae*

Question Number : 103 Question Id : 2106887307 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The number of neutrons, protons and electrons of ${}_{17}\text{Cl}^{37}$ are

Options :

- 1. ✖ 20, 20, 17
- 2. ✖ 17,17,20
- 3. ✔ 20, 17,17
- 4. ✖ 17,17,17

Question Number : 104 Question Id : 2106887308 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The cell organelle involved in the synthesis of phospholipids is

Options :

- 1. ✖ Golgi
- 2. ✖ Chloroplast
- 3. ✖ Lysosomes
- 4. ✔ Endoplasmic reticulum

Question Number : 105 Question Id : 2106887309 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time : 0

The biocontrol agent used against caterpillars is

Options :

1. ✖ *Trichoderma viridae*
2. ✖ *Streptococcus polymixia*
3. ✔ *Bacillus Thuringiensis*
4. ✖ *Saccharomyces cerevisiae*

Question Number : 106 Question Id : 2106887310 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Media that forms foam is

Options :

1. ✖ Natural Media
2. ✖ Synthetic Media
3. ✔ Complex Media
4. ✖ Defined Media

Question Number : 107 Question Id : 2106887311 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The role of calcium in a media is

Options :

1. ✘ Synthesis of phospholipids
2. ✘ Synthesis of Vitamins
3. ✔ Stabilization of cell wall
4. ✘ Stabilization of nucleic acids

Question Number : 108 Question Id : 2106887312 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In pyrethroids, which channel is targeted for imposing the toxicity in pests

Options :

1. ✘ Potassium
2. ✔ Sodium
3. ✘ Calcium

4. ✖ Iron

Question Number : 109 Question Id : 2106887313 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

----- is a major component of a Bordeaux Mixture

Options :

1. ✖ Sodium Chloride

2. ✖ Calcium Chloride

3. ✔ Copper Sulphate

4. ✖ Magnesium Sulphate

Question Number : 110 Question Id : 2106887314 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

----- is a nitrogen fixer in paddy fields

Options :

1. ✖ *Frankia*

2. ✖ *Oscillatoria*

3. ✓ *Azospirillum*

4. ✗ *Rhizobium*

Question Number : 111 Question Id : 2106887315 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The focal lengths of the objective and the eye piece of a compound microscope are 0.95 cm and 5 cm respectively. They are kept at a distance of 20 cm. A large image is formed at a distance of 25 cm. The total magnification is

Options :

1. ✗ 84

2. ✗ 80

3. ✓ 94

4. ✗ 90

Question Number : 112 Question Id : 2106887316 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Resolving power of a microscope can be enhanced by

Options :

1. ✗

Illumination of longer wave length and decreased numerical aperture

2. ✖ Illumination of longer wave length and increased numerical aperture

3. ✖ Illumination of shorter wave length and decreased numerical aperture

4. ✔ Illumination of shorter wave length and increased numerical aperture

Question Number : 113 Question Id : 2106887317 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A plasma membrane is composed of

Options :

1. ✖ A protein, lipid and a cellulose layer

2. ✔ Bimolecular lipid layer surrounded by protein layer

3. ✖ A protein layer between two lipid layers

4. ✖ A lipid layer between two protein layers

Question Number : 114 Question Id : 2106887318 Display Question Number : Yes Is Question

Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The cross of F1 to either of its parents is known as

Options :

1. ✖ Back cross
2. ✔ Test cross
3. ✖ F1 cross
4. ✖ F2 cross

Question Number : 115 Question Id : 2106887319 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Prokaryote and eukaryote cells have a common

Options :

1. ✖ Histone
2. ✖ Cell Division Apparatus
3. ✔ Mitochondria
4. ✖ Genetic code

Question Number : 116 Question Id : 2106887320 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The hydrophobic ends of a phospholipid are

Options :

1. ✖ Polar
2. ✔ Non-Polar
3. ✖ Bi-polar
4. ✖ Neutral

Question Number : 117 Question Id : 2106887321 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

An example of a peripheral membrane protein is

Options :

1. ✖ Insulin receptor
2. ✖ Glycophorin
3. ✖ Integrins

4. ✓ Glycolipid transfer proteins

Question Number : 118 Question Id : 2106887322 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The maximum mass of an atom is concentrated in

Options :

1. ✓ Nucleus

2. ✗ Protons

3. ✗ Neutrons

4. ✗ Electrons

Question Number : 119 Question Id : 2106887323 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Major interactions that stabilize a plasma membrane are

Options :

1. ✗ Hydrophilic interactions

2. ✓

Hydrophobic interactions

3. ✖ Covalent interactions

4. ✖ Ionic interactions

Question Number : 120 Question Id : 2106887324 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Ion carriers are located on

Options :

1. ✖ Nucleus

2. ✖ Cytoplasm

3. ✔ Plasma membrane

4. ✖ Cell wall

Question Number : 121 Question Id : 2106887325 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A cross in which parents differ in a single pair of contrasting character are called

Options :

1. ✓ Monohybrid Cross
2. ✗ Dihybrid Cross
3. ✗ Trihybrid Cross
4. ✗ Tetrahybrid Cross

Question Number : 122 Question Id : 2106887326 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Linkage results in the formation of more

Options :

1. ✗ Dominant phenotype
2. ✗ Wild phenotype
3. ✓ Parental phenotype
4. ✗ Recombinant phenotype

Question Number : 123 Question Id : 2106887327 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

F3 Generation is obtained by

Options :

1. ✖ Selfing of F1
2. ✔ Selfing of F2
3. ✖ Crossing of F1 with F2
4. ✖ Crossing F1 with any one parent

Question Number : 124 Question Id : 2106887328 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In a linear chromosome map, if the distance between four loci are: a-b is 10%, a-d is 3%, b-c is 4%, and a-c is 6%, the frequency of crossover between c and d is

Options :

1. ✔ 3%
2. ✖ 6%

3. ✖ 9%

4. ✖ 12%

Question Number : 125 Question Id : 2106887329 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

As the distance between two genes decreases, linkage will

Options :

1. ✔ Increase

2. ✖ Decrease

3. ✖ Unaffected

4. ✖ Complete

Question Number : 126 Question Id : 2106887330 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Recombination nodules appear in ----- of meiosis

Options :

1. ✖

Leptotene

2. ✖ Zygotene

3. ✔ Pachytene

4. ✖ Diakinesis

Question Number : 127 Question Id : 2106887331 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Spindle fibers are formed and get attached to the chromosomes during

Options :

1. ✖ Prophase

2. ✔ Metaphase

3. ✖ Anaphase

4. ✖ Telophase

Question Number : 128 Question Id : 2106887332 Display Question Number : Yes Is Question

Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Lysosome is formed from which of these cell organelles ?

Options :

1. ✖ Nucleus
2. ✖ Endoplasmic Reticulum
3. ✔ Golgi bodies
4. ✖ Ribosomes

Question Number : 129 Question Id : 2106887333 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Daughter cells receive an exact copy of chromosomes from parent cells in

Options :

1. ✖ Meiosis I
2. ✖ Meiosis II
3. ✔ Mitosis

4. ✖ Interphase

Question Number : 130 Question Id : 2106887334 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Microtubule that pulls the chromosomes towards poles during mitosis is

Options :

1. ✔ Kinetochore

2. ✖ Centriole

3. ✖ Polar

4. ✖ Astra

Question Number : 131 Question Id : 2106887335 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The DNA filaments that are developed inside the nucleus during cell division are

Options :

1. ✖ Spindle Fibers

2. ✔ Chromosomes

3. ✖ Centrioles

4. ✖ Kinetochore

Question Number : 132 Question Id : 2106887336 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

----- is required for the chromosomal condensation while the cell enters mitosis

Options :

1. ✖ Condensin

2. ✖ Histone

3. ✔ Cohesin

4. ✖ Topoisomerase

Question Number : 133 Question Id : 2106887337 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of these histones binds to a linker DNA

Options :

1. ✔

H1

2. ✖ H2A

3. ✖ H2B

4. ✖ H3

Question Number : 134 Question Id : 2106887338 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The DNA in a cell remains uncondensed throughout

Options :

1. ✔ Interphase

2. ✖ Prophase

3. ✖ Metaphase

4. ✖ Telophase

Question Number : 135 Question Id : 2106887339 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time : 0

In male humans, the gene for colour blindness is located on

Options :

1. ✓ X-Chromosome

2. ✗ Y-Chromosome

3. ✗ chromosome 12

4. ✗ Chromosome 14

Question Number : 136 Question Id : 2106887340 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A completely expressed allele is known as

Options :

1. ✓ Dominant

2. ✗ Recessive

3. ✗ Homozygous

4.

✖ Heterozygous

Question Number : 137 Question Id : 2106887341 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The genotype of a person suffering from Klinefelter's syndrome is

Options :

1. ✖ 44+XXX

2. ✖ 42+XXX

3. ✔ 44+XXY

4. ✖ 42+XYY

Question Number : 138 Question Id : 2106887342 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A female with XXXX chromosome will have ----- number of Barr bodies

Options :

1. ✖ One

2. ✖ Two

3. ✔ Three

4. ✖ Four

Question Number : 139 Question Id : 2106887343 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Sex determination in *Drosophila* is by

Options :

1. ✖ Autosomes

2. ✔ X Chromosome and Autosomes

3. ✖ Y Chromosomes

4. ✖ Y Chromosome and Autosomes

Question Number : 140 Question Id : 2106887344 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Genes that are present in the homologous region of X and Y chromosomes are known as

Options :

1. ✖ Autosomal
2. ✖ Sex-linked
3. ✔ Partially Sex linked
4. ✖ Unlinked

Question Number : 141 Question Id : 2106887345 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If a person is colour blind, the probability of his son getting colour blindness is

Options :

1. ✔ 0
2. ✖ $1/4$
3. ✖ $1/2$
4. ✖ 1

Question Number : 142 Question Id : 2106887346 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A Y-linked inheritance is

Options :

1. ✖ Criss-Cross

2. ✖ Loop

3. ✔ Straight

4. ✖ Jumping

Question Number : 143 Question Id : 2106887347 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Genes that are located only on Y-chromosome are known as

Options :

1. ✖ Lethal Genes

2. ✖ Duplicate genes

3. ✖ Plasma genes

4. ✓ Holandric genes

Question Number : 144 Question Id : 2106887348 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The condition in which a single gene controls more than one completely un-related phenotypic trait is known as

Options :

1. ✗ Polygenes

2. ✗ Multiple allelism

3. ✗ Multiple genes

4. ✓ Pleiotropism

Question Number : 145 Question Id : 2106887349 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Down's syndrome is an example of

Options :

1. ✗ Monosomy

2. ✓ Trisomy

3. ✖ Triploidy

4. ✖ Eupolyploidy

Question Number : 146 Question Id : 2106887350 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

% similarity of each strain to every other strain is calculated by

Options :

1. ✖ DNA homology

2. ✖ Genetic relatedness

3. ✔ Numerical Taxonomy

4. ✖ Intuitive method

Question Number : 147 Question Id : 2106887351 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

According to Bergey's Manual of Systematic Bacteriology, prokaryotes that lack a cell wall belong to

Options :

1. ✖

Gracilicutes

2. ✖ Firmicutes

3. ✔ Tenericutes

4. ✖ Mendosicutes

Question Number : 148 Question Id : 2106887352 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The type of bacteria that uses carbon dioxide as a sole source for their growth are

Options :

1. ✖ Lithotrophs

2. ✔ Autotrophs

3. ✖ Heterotrophs

4. ✖ Organotrophs

Question Number : 149 Question Id : 2106887353 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time : 0

The generation time of *E. coli* is

Options :

1. ✖ 13 minutes
2. ✔ 20 minutes
3. ✖ 45 Minutes
4. ✖ 3 minutes

Question Number : 150 Question Id : 2106887354 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

During exponential phase the microbial growth rate is

Options :

1. ✖ Similar to generation time
2. ✔ Reciprocal of generation time
3. ✖ Double the generation time
4. ✖ Double the population

Question Number : 151 Question Id : 2106887355 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Roll tube technique is used for the isolation of

Options :

1. ✖ Aerobes
2. ✖ Anaerobes
3. ✖ Facultatives
4. ✔ Stringent anaerobes

Question Number : 152 Question Id : 2106887356 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The unit of influent flow rate is

Options :

1. ✖ md
2. ✖ m/d
3. ✖ m²/d

4. ✓ m^3/d

Question Number : 153 Question Id : 2106887357 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The reverse of hydraulic retention time is

Options :

1. ✗ Chemical rate

2. ✓ Dilution rate

3. ✗ Filtration rate

4. ✗ Sedimentation rate

Question Number : 154 Question Id : 2106887358 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

In which of the bioreactors particles are not immersed in liquid

Options :

1. ✗ Air-lift

2. ✖ Stirred vessel

3. ✖ Packed-bed

4. ✔ Trickle –bed

Question Number : 155 Question Id : 2106887359 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

An agent that prevent the growth of bacteria is known as

Options :

1. ✖ Antibiotic

2. ✖ Bactericide

3. ✔ Bacteriostatic

4. ✖ Antimicrobial

Question Number : 156 Question Id : 2106887360 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The pressure at which the autoclave operated is

Options :

1. ✖ 5 Psi
2. ✖ 10 Psi
3. ✔ 15 Psi
4. ✖ 20 Psi

Question Number : 157 Question Id : 2106887361 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If the reactions $A \rightarrow B$ and $A \rightarrow C$ are represented as $(-r_A) = k_1 C_A^3$ and $(-r_A) = k_2 C_A^6$ of activation energies 130 and 150 J/ mol, then which of the following is the right selection of process parameters to increase selectivity of B?

Options :

1. ✖ CSTR, High temperature
2. ✖ PFR, High temperature
3. ✔ CSTR, Low temperature
4. ✖ PFR, Low temperature

Question Number : 158 Question Id : 2106887362 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which form of DNA structure is described by Watson-Crick model

Options :

1. ✓ B Form

2. ✗ Z form

3. ✗ A-Form

4. ✗ D- Form

Question Number : 159 Question Id : 2106887363 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The stability and the formation of a G-quadruplex depends on

Options :

1. ✓ Monovalent cation

2. ✗ Bivalent cation

3. ✗ Divalent cation

4. ✖ Pentavalent cation

Question Number : 160 Question Id : 2106887364 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which form of structure is adapted by RNA

Options :

1. ✔ B Form

2. ✖ Z form

3. ✖ A-Form

4. ✖ D- Form

Question Number : 161 Question Id : 2106887365 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

RNA molecules that carry the genetic information copied from DNA are

Options :

1. ✖ miRNA

2. ✖

rRNA

3. ✖ tRNA

4. ✔ mRNA

Question Number : 162 Question Id : 2106887366 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following is not a base of RNA ?

Options :

1. ✖ Adenine

2. ✖ Guanine

3. ✔ Thymine

4. ✖ Cytosine

Question Number : 163 Question Id : 2106887367 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which enzyme will remove the supercoiling in DNA ahead of the replication fork?

Options :

1. ✖ DNA Polymerase
2. ✖ DNA gyrase
3. ✔ Topoisomerase
4. ✖ Helicase

Question Number : 164 Question Id : 2106887368 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following is the principal replication enzyme of *E coli* ?

Options :

1. ✖ DNA Polymerase I
2. ✖ DNA Polymerase II
3. ✔ DNA Polymerase III
4. ✖ DNA polymerase IV

Question Number : 165 Question Id : 2106887369 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Association of DNA and histone protein is mediated by

Options :

1. ✖ Covalent bonding
2. ✔ Hydrogen bonding
3. ✖ Ionic bonding
4. ✖ Vander Waals interactions

Question Number : 166 Question Id : 2106887370 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The equation demonstrating a DNA renaturation reaction is

Options :

1. ✖ Sec 60
2. ✔ Cot1/2
3. ✖ Tan 30

4. ✖ Cot 40

Question Number : 167 Question Id : 2106887371 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The location of microsatellite repeats on a chromosome is

Options :

1. ✖ Centromere

2. ✖ Telomere

3. ✖ Acromere

4. ✔ Dispersed throughout the chromosome

Question Number : 168 Question Id : 2106887372 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A nucleosome is composed of

Options :

1. ✖ DNA and Histone protein

2. ✓ DNA, Histone protein and linker H1

3. ✗ RNA and Histone protein

4. ✗ RNA, Histone protein and linker H1

Question Number : 169 Question Id : 2106887373 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The number of stop codons in a genetic code is

Options :

1. ✗ 1

2. ✗ 2

3. ✓ 3

4. ✗ 4

Question Number : 170 Question Id : 2106887374 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Loops in lampbrush chromosomes represent the sites of

Options :

1. ✖ Replication
2. ✔ Transcription
3. ✖ Linkage
4. ✖ Crossingover

Question Number : 171 Question Id : 2106887375 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Human ABO blood group is an example of

Options :

1. ✖ Dominance
2. ✖ Recessiveness
3. ✔ Co-dominance
4. ✖ Segregation

Question Number : 172 Question Id : 2106887376 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

If the percentage of adenine is 30 in a DNA molecule, then what would be the guanine, thymine and cytosine

Options :

- 1. ✖ T:20 C:25 G: 25
- 2. ✖ T:25 C:20 G:25
- 3. ✖ T:20 C:20 G:30
- 4. ✔ T:30 C:20 G:20

Question Number : 173 Question Id : 2106887377 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Alkaptonuria and Albinism occurs through

Options :

- 1. ✖ Autosomal Dominant inheritance
- 2. ✔ Autosomal Recessive inheritance
- 3. ✖ X-Chromosome Inheritance

4. ✖ Y-Chromosome inheritance

Question Number : 174 Question Id : 2106887378 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which among the following is correct ?

Options :

1. ✖ Down syndrome: 44 Autosomes + XO

2. ✔ Klinefelter's syndrome :44 Autosomes + XXY

3. ✖ Erythroblastosis fetalis: X linked

4. ✖ Colour blindness: Y linked

Question Number : 175 Question Id : 2106887379 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following RNAs are not required for the protein synthesis ?

Options :

1. ✔ siRNA

2. ✖ mRNA

3. ✖ tRNA

4. ✖ rRNA

Question Number : 176 Question Id : 2106887380 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

What is the size of a prokaryote ribosome ?

Options :

1. ✖ 40S

2. ✖ 60S

3. ✔ 70S

4. ✖ 80S

Question Number : 177 Question Id : 2106887381 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of these is a start codon in prokaryotes ?

Options :

1. ✖ AUG

2. ✔ GUG

3. ✖ UAA

4. ✖ UAG

Question Number : 178 Question Id : 2106887382 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Viruses that infect bacteria are known as

Options :

1. ✖ Archaea

2. ✔ Phages

3. ✖ phases

4. ✖ pUC

Question Number : 179 Question Id : 2106887383 Display Question Number : Yes Is Question

Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The process by which a foreign DNA is introduced into a bacterial cell is Known as

Options :

1. ✓ Transformation

2. ✗ Transduction

3. ✗ Digestion

4. ✗ Amplification

Question Number : 180 Question Id : 2106887384 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The organism that can transfer t-DNA into the plants is

Options :

1. ✗ *E.coli*

2. ✓ *Agrobacterium tumifaciens*

3. ✗ *Aspergillus niger*

Trichoderma viridae

4. ✖

Question Number : 181 Question Id : 2106887385 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Expression vectors differ from a cloning vector in

Options :

1. ✖ Origin of replication

2. ✖ Suitable marker genes

3. ✖ Restriction site

4. ✔ regulatory elements

Question Number : 182 Question Id : 2106887386 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

cDNA libraries are produced from

Options :

1. ✔ mRNA

2. ✖ tRNA

3. ✖ rRNA

4. ✖ miRNA

Question Number : 183 Question Id : 2106887387 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which enzyme is used in the PCR amplification of DNA

Options :

1. ✔ DNA polymerase

2. ✖ Hexonuclease

3. ✖ RNA Polymerase

4. ✖ Endonuclease

Question Number : 184 Question Id : 2106887388 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which tissue culture technique is applied for producing virus free plants ?

Options :

1. ✖ Embryo culture
2. ✖ Ovule culture
3. ✔ Meristem culture
4. ✖ Anther culture

Question Number : 185 Question Id : 2106887389 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

What is protoplast ?

Options :

1. ✖ Cell wall and Plasma membrane of a cell
2. ✖ Cytoplasm with cell wall
3. ✔ Cell without a cell wall
4. ✖ cytoplasm without plasma membrane

Question Number : 186 Question Id : 2106887390 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The function of cytokines in plant tissue culture is

Options :

1. ✓ Adventitious shoot formation
2. ✗ Somatic embryo induction
3. ✗ Adventitious root formation
4. ✗ Shoot elongation

Question Number : 187 Question Id : 2106887391 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Callus formation is induced by

Options :

1. ✓ Auxins
2. ✗ Cytokines
3. ✗

Gibberellin

4. ✖ Ethylene

Question Number : 188 Question Id : 2106887392 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Fusogenic agent used for the protoplast culture is

Options :

1. ✔ PEG

2. ✖ SDS

3. ✖ PUC

4. ✖ Salicylic acid

Question Number : 189 Question Id : 2106887393 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The culture prepared by inoculating from the tissue of an organism to the culture media is

Options :

1. ✔ Primary cell culture

Secondary culture

2. ✖

Cell lines

3. ✖

Transformed cell culture

4. ✖

Question Number : 190 Question Id : 2106887394 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The ratio of CO₂: O₂ used in the cell culture system is

Options :

1:5

1. ✖

1:13

2. ✖

1:19

3. ✔

1:23

4. ✖

Question Number : 191 Question Id : 2106887395 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time : 0

Protein produced from transgenic sheep used in the replacement therapy for individuals suffering with emphysema is

Options :

1. ✖ Plasminogen activator
2. ✔ α -antitrypsin
3. ✖ Casein
4. ✖ Amyloid precursor proteins

Question Number : 192 Question Id : 2106887396 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Which of the following is a biological method for gene transfer ?

Options :

1. ✖ Microinjection
2. ✖ Electroporation
3. ✖ Particle bombardment
4. ✔

Baculoviral vector system

Question Number : 193 Question Id : 2106887397 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

A cell line is a

Options :

1. ✖ multilayer culture
2. ✖ Transformed cells
3. ✖ Multiple growth of cells
4. ✔ Sub-culture of a primary culture

Question Number : 194 Question Id : 2106887398 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Under favourable conditions and chemically defined growth media, protoplasts can grow into

Options :

1. ✖ Organ

2.

✓ Callus

3. ✗ Root

4. ✗ Shoot

Question Number : 195 Question Id : 2106887399 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The process of de-differentiation in a cell culture can give rise to

Options :

1. ✗ Carcinoma cells

2. ✗ Single protoplasts

3. ✓ Induced-pluripotent stem cells

4. ✗ Fused protoplasts

Question Number : 196 Question Id : 2106887400 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Metabolites that are formed during growth phase as an outcome of energy metabolism are known as

Options :

1. ✓ Primary metabolites
2. ✗ Secondary metabolites
3. ✗ Tertiary metabolites
4. ✗ Quaternary metabolites

Question Number : 197 Question Id : 2106887401 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

The primary storage of food in animals is through

Options :

1. ✗ Glucose
2. ✗ Galactose
3. ✗ Glycogen
4. ✓ Fats


Question Number : 198 Question Id : 2106887402 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Apoenzyme is a

Options :

1. ✓ protein portion of an enzyme
2. ✗ Non-protein portion of an enzyme
3. ✗ complete biologically active conjugated enzyme
4. ✗ prosthetic group

Question Number : 199 Question Id : 2106887403 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0



Options :

1. ✓ Hydrolases
2. ✗ Aldolase

3. ✖ Peroxidases

4. ✖ Pectinases

Question Number : 200 Question Id : 2106887404 Display Question Number : Yes Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Enzymes catalyse biochemical reactions by altering the following

Options :

1. ✖ The enthalpy of formation ΔH

2. ✖ The equilibrium constant K_e

3. ✖ Change in the Gibb's free energy ΔG

4. ✔ The activation energy E_a