

WARNING		Any malpractice or any attempt to commit any kind of malpractice in the Examination will DISQUALIFY THE CANDIDATE .	
PAPER – II MATHEMATICS-2020			
Version Code	B1	Question Booklet Serial Number :	9126960
Time: 150 Minutes		Number of Questions: 120	Maximum Marks: 480
Name of the Candidate			
Roll Number			
Signature of the Candidate			
INSTRUCTIONS TO CANDIDATES			
1. Please ensure that the VERSION CODE shown at the top of this Question Booklet is same as that shown in the Admit Card issued to you. If you have received a Question Booklet with a different Version Code please get it replaced with a Question Booklet with the same Version Code as in the Admit Card from the Invigilator. THIS IS VERY IMPORTANT.			
2. Please fill the items such as Name, Roll Number and Signature in the columns given above. Please also write Question Booklet Serial Number given at the top of this page against item 3 in the OMR Answer Sheet.			
3. This Question Booklet contains 120 questions. For each question five answers are suggested and given against (A), (B), (C), (D) and (E) of which only one will be the 'Most Appropriate Answer.' Mark the bubble containing the letter corresponding to the 'Most Appropriate Answer' in the OMR Answer Sheet, by using either Blue or Black Ball Point Pen only.			
4. Negative Marking: In order to discourage wild guessing the score will be subjected to penalization formula based on the number of right answers actually marked and the number of wrong answer marked. Each correct answer will be awarded FOUR marks. ONE mark will be deducted for each incorrect answer. More than one answer marked against a question will be deemed as incorrect answer and will be negatively marked.			
5. Please read the instructions in the OMR Answer Sheet for marking the answers. Candidates are advised to strictly follow the instruction contained in the OMR Answer Sheet.			
IMMEDIATELY AFTER OPENING THE QUESTION BOOKLET, THE CANDIDATE SHOULD VERIFY WHETHER THE QUESTION BOOKLET CONTAINS ALL THE 120 QUESTIONS IN SERIAL ORDER. IF NOT, REQUEST FOR REPLACEMENT.			
DO NOT OPEN THE SEAL UNTIL THE INVIGILATOR ASKS YOU TO DO SO.			

SEAL

Name of the Candidate: _____
 Roll Number: _____
 Date: _____
 Time: _____
 Signature of the Candidate: _____
 Head of the Institution: _____
 Name of the Institution: _____
 Address: _____
 City: _____
 State: _____
 Pin Code: _____
 District: _____
 Country: _____
 Phone Number: _____
 Email Address: _____
 Website: _____
 Social Media: _____
 Other: _____
 DO NOT WRITE ANYTHING ON THE BACK OF THE QUESTION BOOKLET.

BLANK PAGE

PLEASE ENSURE THAT THIS QUESTION BOOKLET CONTAINS
120 QUESTIONS SERIALLY NUMBERED FROM 1 TO 120
PRINTED PAGES 32.

1. The domain of the function f given by $f(x) = \sqrt{x-1}$ is
(A) $(-\infty, \infty)$ (B) $(1, \infty)$ (C) $[1, \infty)$ (D) $[0, \infty)$ (E) $(0, \infty)$
2. Let $f(x) = -2x^2 + 1$ and $g(x) = 4x - 3$, then $(g \circ f)(-1)$ is equal to
(A) 9 (B) -9 (C) 7 (D) -7 (E) -8
3. Let A and B be finite sets such that $n(A - B) = 18$, $n(A \cap B) = 25$ and $n(A \cup B) = 70$.
Then $n(B)$ is equal to
(A) 52 (B) 25 (C) 27 (D) 43 (E) 45
4. In a group of 100 persons, 80 people can speak Malayalam and 60 can speak English.
Then the number of people who speak English only is
(A) 40 (B) 30 (C) 20 (D) 25 (E) 35

Space for rough work

5. If $*$ is a binary operation defined by $a * b = \frac{a}{b} + \frac{b}{a} + \frac{1}{ab}$ for positive integers a and b , then $2 * 5$ is equal to
 (A) 4 (B) 3 (C) 2 (D) 1 (E) 5
6. If $A = \{1, 2, 3, 4, 5\}$ and $B = \{2, 4, 6\}$, then $A - B =$
 (A) $\{1, 3, 5, 6\}$ (B) $\{0, 1, 3, 5, 6\}$ (C) $\{1, 3, 5\}$
 (D) $\{1, 2, 3, 4, 5, 6\}$ (E) $\{2, 4\}$
7. Let $A = \{2, 3, 4, 5\}$, $B = \{36, 45, 49, 60, 77, 90\}$ and let R be the relation 'is factor of' from A to B . Then the range of R is the set
 (A) $\{60\}$ (B) $\{36, 45, 60, 90\}$ (C) $\{49, 77\}$
 (D) $\{49, 60, 77\}$ (E) $\{36, 45, 49, 60, 77, 90\}$
8. The real part of $e^{(3+4i)x}$ is
 (A) e^{3x} (B) $\cos 7x$ (C) $e^{3x} \cos 4x$
 (D) $e^{3x} \sin 4x$ (E) 0
9. If $z = x - iy$ and $z^{1/3} = p + iq$, then $\frac{1}{p^2 + q^2} \left(\frac{x}{p} + \frac{y}{q} \right)$ is equal to
 (A) -2 (B) -1 (C) 1 (D) 2 (E) 0

Space for rough work

10. Let $z = x + iy$ be a complex number such that $|z + i| = 2$. Then the locus of z is a circle whose centre and radius are
- (A) $(0, -1); 2$ (B) $(0, 2); 2$ (C) $(1, -1); 2$
 (D) $(0, -1); \sqrt{3}$ (E) $(0, 2); \sqrt{3}$
11. If $2 + i$ is a root of $x^2 - 4x + c = 0$, where c is a real number, then the value of c is
- (A) 2 (B) 3 (C) 4 (D) 5 (E) 0
12. Let z_1 and z_2 be complex numbers satisfying $|z_1| = |z_2| = 2$ and $|z_1 + z_2| = 3$.
 Then $\left| \frac{1}{z_1} + \frac{1}{z_2} \right| =$
- (A) $\frac{3}{2}$ (B) 2 (C) $\frac{3}{4}$ (D) $\frac{1}{2}$ (E) 4
13. The principal argument of the complex number $z = \frac{1 + \sin \pi - i \cos \pi}{1 + \sin \pi + i \cos \pi}$ is
- (A) $\frac{\pi}{3}$ (B) $\frac{\pi}{6}$ (C) $\frac{\pi}{5}$ (D) $\frac{\pi}{2}$ (E) $\frac{\pi}{4}$

Space for rough work

14. If $z_1 = 2 + 3i$ and $z_2 = 3 + 2i$, then $|z_1 + z_2|$ is equal to
(A) 50 (B) 10 (C) $5\sqrt{2}$ (D) 25 (E) $2\sqrt{5}$
15. $\frac{10i}{1+2i}$ is equal to
(A) $-2i$ (B) $2i$ (C) $-4 + 2i$ (D) $4 + 2i$ (E) $6i$
16. The value of $\sum_{k=1}^{10} (3k^2 + 2k - 1)$ is
(A) 1120 (B) 1200 (C) 1230 (D) 1265 (E) 1255
17. The numbers $a_1, a_2, a_3, \dots$ form an arithmetic sequence with $a_1 \neq a_2$. The three numbers a_1, a_2 and a_6 form a geometric sequence in that order. Then the common difference of the arithmetic sequence is
(A) a_1 (B) $2a_1$ (C) $3a_1$ (D) $4a_1$ (E) $5a_1$
18. In an arithmetic sequence, the sum of first and third terms is 6 and the sum of second and fourth terms is 20. Then the 11th term is
(A) 67 (B) 62 (C) 57 (D) 73 (E) 66

Space for rough work

19. In an A.P., the first term is 3 and the last term is 17. The sum of all the terms in the sequence is 70. Then the number of terms in the arithmetic sequence is
(A) 7 (B) 5 (C) 9 (D) 6 (E) 8
20. Consider the set of all positive rational numbers that are less than 1 and that have denominators as 30 in their lowest terms. Their sum is equal to
(A) 1 (B) 2 (C) 3 (D) 4 (E) 5
21. If p , q and 23 is an increasing arithmetic sequence and p and q are prime numbers, then $p + q =$
(A) 22 (B) 24 (C) 26 (D) 28 (E) 30
22. The 5th and 7th terms of a G.P. are 12 and 48 respectively. Then the 9th term is
(A) 162 (B) 96 (C) 192 (D) 144 (E) 182
23. The number of positive integers less than 1000 having only odd digits is
(A) 155 (B) 177 (C) 55 (D) 205 (E) 85

Space for rough work

24. Five points are marked on a circle. The number of distinct polygons of three or more sides can be drawn using some (or all) of the five points as vertices is
 (A) 10 (B) 12 (C) 14 (D) 16 (E) 18
25. The middle term in the expansion of $\left(1 + \frac{1}{5}\right)^{20}$ is
 (A) $\left(\frac{1}{5}\right)^{10}$ (B) $\left(\frac{1}{5}\right)^{11}$ (C) ${}^{20}C_{11}\left(\frac{1}{5}\right)^{11}$ (D) ${}^{20}C_9\left(\frac{1}{5}\right)^9$ (E) ${}^{20}C_{10}\left(\frac{1}{5}\right)^{10}$
26. ${}^{11}C_0 + {}^{11}C_1 + {}^{11}C_2 + {}^{11}C_3 + {}^{11}C_4 + {}^{11}C_5 =$
 (A) 2^6 (B) 2^8 (C) 2^{10} (D) 2^{11} (E) 2^9
27. If ${}^nP_r = 840$ and ${}^nC_r = 35$, then the value of r is equal to
 (A) 2 (B) 4 (C) 6 (D) 3 (E) 5
28. The sum of the coefficients in the expansion of $(1 + 2x - x^2)^{20}$ is
 (A) 2^{20} (B) 2^{21} (C) 2^{19} (D) 2^{40} (E) 2

Space for rough work

29. The number of ways a committee of 4 people can be chosen from a panel of 10 people is

- (A) 315 (B) 240 (C) 210 (D) 720 (E) 120

30. If $A = \begin{pmatrix} 6 & 2 \\ 7 & -5 \end{pmatrix}$ and $A - B = \begin{pmatrix} -2 & 1 \\ 4 & -9 \end{pmatrix}$, then $B =$

- (A) $\begin{pmatrix} -8 & -1 \\ 3 & 4 \end{pmatrix}$ (B) $\begin{pmatrix} 8 & 1 \\ -3 & -4 \end{pmatrix}$ (C) $\begin{pmatrix} 4 & 3 \\ 11 & -14 \end{pmatrix}$ (D) $\begin{pmatrix} 8 & 1 \\ 3 & 4 \end{pmatrix}$ (E) $\begin{pmatrix} 4 & 1 \\ 3 & 2 \end{pmatrix}$

31. The value of the determinant $\begin{vmatrix} bc & ca & ab \\ a^3 & b^3 & c^3 \\ \frac{1}{a} & \frac{1}{b} & \frac{1}{c} \end{vmatrix}$ is

- (A) $a^5 - 1$ (B) $a^2bc + ab^2c + abc^2$ (C) $ab(a+b+c)$
(D) $a^4b^4c^4(a+b+c)$ (E) 0

32. If the matrix $\begin{bmatrix} 1 & 2 & -1 \\ -3 & 4 & k \\ -4 & 2 & 6 \end{bmatrix}$ is singular, then the value of k is equal to

- (A) 3 (B) 4 (C) 5 (D) 6 (E) 7

Space for rough work

33. If $\begin{bmatrix} -1 & 3 \\ 4 & -5 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 7 \end{bmatrix} = \begin{bmatrix} -1 & 19 \\ \alpha & -27 \\ 0 & 14 \end{bmatrix}$, then the value of α is

(A) 5 (B) 4 (C) 7 (D) -14 (E) -5

34. If $A^{-1} = \frac{1}{11} \begin{pmatrix} -3 & 4 \\ 5 & -3 \end{pmatrix}$, then $A =$

(A) $\frac{-1}{11} \begin{pmatrix} 3 & 4 \\ 5 & 3 \end{pmatrix}$ (B) $\frac{1}{11} \begin{pmatrix} 3 & 4 \\ 5 & 3 \end{pmatrix}$ (C) $\begin{pmatrix} 3 & -4 \\ -5 & 3 \end{pmatrix}$

(D) $\begin{pmatrix} 3 & 4 \\ 5 & 3 \end{pmatrix}$ (E) $\begin{pmatrix} -3 & 4 \\ 5 & -3 \end{pmatrix}$

35. The system of equations

$$x + y + 2z = 4$$

$$3x + 3y + 6z = 17$$

$$5x - 3y + 2z = 27$$

has

- (A) no solution (B) finitely many solutions
 (C) infinitely many solutions (D) unique and trivial solution
 (E) unique and non-trivial solution

Space for rough work

36. The smallest prime number satisfying the inequality $\frac{2n-3}{3} \geq \frac{n-1}{6} + 1$ is
 (A) 2 (B) 3 (C) 5 (D) 7 (E) 11
37. The number of integers satisfying the inequality $|n^2 - 100| < 50$ is
 (A) 5 (B) 6 (C) 12 (D) 8 (E) 10
38. The solution set of the rational inequality $\frac{x+9}{x-6} \leq 0$ is
 (A) $(-\infty, 9) \cup (6, \infty)$ (B) $(-\infty, 9] \cup (6, \infty)$ (C) $(-\infty, 9] \cup [6, \infty)$
 (D) $[-9, 6)$ (E) $(-9, 6]$
39. Which of the following sentences is/are statement(s)?
 (i) 10 is less than 5.
 (ii) All rational numbers are real numbers.
 (iii) Today is a sunny day.
 (A) (i), (ii) and (iii) (B) (i) and (ii) only (C) (i) and (iii) only
 (D) (ii) and (iii) only (E) (i) only

Space for rough work

