



### General Instructions

- (i) This question paper contains 200 questions. All questions are compulsory.
- (ii) It comprises 200 single-correct multiple-choice questions.
- (iii) These questions follow the NEET PG examination pattern.
- (iv) Each correct answer is awarded +4 marks, with -1 for an incorrect answer.
- (v) The actual NEET paper has 180 questions to be attempted in 3 hours 20 minutes.
- (vi) Use of calculators is not permitted.
- (vii) Attempt every question; detailed solutions are provided in the companion solutions booklet.

1. Which muscle is inserted into the floor of the intertubercular sulcus of the humerus?

- (A) Latissimus dorsi
- (B) Teres major
- (C) Pectoralis major
- (D) Deltoid

• • •

2. All are derived from ectoderm except -

- (A) Hypophysis
- (B) Retina
- (C) Spinal cord
- (D) Adrenal cortex

• • •

3. Cytotrophoblast invades which structure?

- (A) Decidua parietalis
- (B) Decidua basalis
- (C) Decidua capsularis
- (D) None

• • •

**4. Inferior parathyroid develops from which pharyngeal arch?**

- (A) 1st
- (B) 2nd
- (C) 3rd
- (D) 4th

• • •

**5. Multi-unit smooth muscle is present at all of the following except?**

- (A) Blood vessels
- (B) Iris
- (C) Gut
- (D) Ductus deferens

• • •

**6. A patient is found to have a melanoma originating in the skin of the left forearm. After removal of the tumor from the forearm, all axillary lymph nodes lateral to the medial edge of the pectoralis minor muscle are removed. Which axillary nodes would NOT be removed?**

- (A) Apical lymph nodes
- (B) Central lymph nodes
- (C) Lateral lymph nodes
- (D) Pectoral lymph nodes

• • •

7. Which structure(s) passes behind the inguinal ligament?

- (A) Femoral branch of genitofemoral nerve
- (B) Femoral vein
- (C) Psoas major
- (D) All

• • •

8. An inferior vena cava (IVC) filter is used for all of the following purposes except:

- (A) To reduce symptoms
- (B) Negligible size of emboli
- (C) To prolong life
- (D) To prevent progression of native blood vessel disease

• • •

9. The surface marking of the oblique fissure of the lung includes all of the following except:

- (A) T3 spine
- (B) 5th rib
- (C) 7th rib
- (D) 6th costal cartilage

• • •

10. Which of the following structures is seen passing through the cavernous sinus?

- (A) Maxillary division of the V nerve
- (B) Mandibular division of the V nerve
- (C) Internal carotid artery
- (D) Facial nerve

• • •

11. Renal vein thrombosis is associated with all of the following except:

- (A) Trauma
- (B) Sick cell anemia
- (C) Nephrotic syndrome
- (D) Dehydration

• • •

**12.** Regarding celiac plexus block, all of the following are true except:

- (A) Relieves pain from gastric malignancy
- (B) Causes hypotension
- (C) Can be used to provide anesthesia for intra-abdominal surgery
- (D) Can be given only by the retrocruel (classic) approach

• • •

**13.** All of the following are branches of the inferior mesenteric artery except:

- (A) Left colic artery
- (B) Sigmoidal artery
- (C) Middle rectal artery
- (D) Superior rectal artery

• • •

**14.** The ophthalmic artery is a branch of which of the following?

- (A) Cavernous part of the internal carotid artery (ICA)
- (B) Cerebral part of the internal carotid artery (ICA)
- (C) Middle cerebral artery (MCA)
- (D) Facial artery

• • •

**15.** Which of the following is not a part of the ethmoid bone?

- (A) Agger nasi
- (B) Crista galli
- (C) Uncinate process
- (D) Inferior turbinate

• • •

**16.** Which of the following laryngeal cartilages is hyaline?

- (A) Epiglottis
- (B) Corniculate
- (C) Cricoid
- (D) Cuneiform

• • •

**17.** The thyrocervical trunk is a branch of which part of the subclavian artery?

- (A) 1st
- (B) 2nd
- (C) 3rd
- (D) None

• • •

**18.** All of the following are lateral branches of the abdominal aorta, EXCEPT:

- (A) Right testicular artery
- (B) Left renal artery
- (C) Inferior mesenteric artery
- (D) Middle suprarenal artery

• • •

**19.** What is the difference between the amount of O<sub>2</sub> inspired (taken up) and the CO<sub>2</sub> expired per minute?

- (A) 20 ml/min
- (B) 50 ml/min
- (C) 75 ml/min
- (D) 100 ml/min

• • •

**20.** What is the approximate daily volume of pancreatic secretion?

- (A) 1.5 L
- (B) 2.5 L
- (C) 5.0 L
- (D) 10 L

• • •

**21.** Einthoven's law is best expressed as:

- (A)  $I + III = II$
- (B)  $I - III = II$
- (C)  $I + II + III = 0$
- (D)  $I + III = aVL$

• • •

**22.** Normal interstitial fluid hydrostatic pressure is?

- (A) 10 to 15 mmHg
- (B) -5 to 0 mmHg
- (C) 20 to 30 mmHg
- (D) -10 to -20 mmHg

• • •

**23.** CSF to plasma glucose ratio is?

- (A) 0.2 - 0.4
- (B) 0.6 - 0.8
- (C) 1.2 - 1.6
- (D) 1.6 - 2.2

• • •

**24.** The distance by which two touch stimuli must be separated to be perceived as two separate stimuli is greatest at?

- (A) The lips
- (B) The palm of the hand
- (C) The back over the scapula
- (D) The dorsum of the hand

• • •

**25.** The resting membrane potential in a nerve fibre:

- (A) Is equal to the potential of ventricular muscle fibre
- (B) Can be measured by surface electrodes
- (C) Increases as extracellular  $K^+$  increases
- (D) Depends upon  $K^+$  equilibrium

• • •

**26.** Which of the following is NOT TRUE about CSF?

- (A) Removal of CSF during dural tap causes intense intracranial headache
- (B) Normally contains no neutrophils
- (C) Formed by arachnoid villi within the ventricles
- (D) pH is less than that of plasma

• • •

**27.** The cephalic phase of gastric secretion occurs?

- (A) On food entering the stomach
- (B) On food entering the intestine
- (C) On seeing food
- (D) On stress

• • •

**28.** What is the effect of acetylcholine on the lower oesophageal sphincter (LES)?

- (A) Contraction
- (B) Relaxation
- (C) No effect
- (D) Contraction followed by relaxation

• • •

**29.** Major adrenal androgen is?

- (A) Testosterone
- (B) 11-hydroxy derivative of androstenedione
- (C) 17-ketosteroid dehydroepiandrosterone
- (D) Cortisol

• • •

**30.** Growth hormone level is decreased in?

- (A) Hypoglycemia
- (B) Fasting
- (C) Sleep
- (D) Exercise

• • •

**31.** Glucose is transported in renal tubular cells by?

- (A) K symport
- (B) K antiport
- (C) Na antiport
- (D) Na cotransport

• • •

**32.** Myosin filament has a fixed length of?

- (A) 0.16 nm
- (B) 1.6 micrometers
- (C) 16 nm
- (D) 1.6 mm

• • •

**33.** Glucose-6-phosphate dehydrogenase needs?

- (A) NAD
- (B) NADP
- (C) FAD
- (D) FMN

• • •

**34.**  $K_m$  value is defined as?

- (A) Substrate concentration at  $V_{max}/2$
- (B) Substrate concentration of twice  $V_{max}$
- (C) Substrate concentration of thrice  $V_{max}$
- (D) Substrate concentration of one third  $V_{max}$

• • •

**35.** In the electron transport chain, cyanide inhibits?

- (A) Complex I
- (B) Cytochrome C oxidase
- (C) Complex IV
- (D) Complex III

• • •

**36.** Enzymes of glycolysis are found in:

- (A) Cytosol
- (B) Cell membrane
- (C) Mitochondria
- (D) Ribosomes

• • •

**37.** Rate limiting step in TCA cycle is catalyzed by:

- (A) Alpha-ketoglutarate synthase
- (B) Fumarase
- (C) Aconitase
- (D) Thiokinase

• • •

**38.** Glycogen storage disease which presents as a lysosomal storage disease is:

- (A) Von Gierke's disease
- (B) Pompe's disease
- (C) McArdle's disease
- (D) Andersen's disease

• • •

**39.** Mutation in GLUT-2 causes:

- (A) Dandy-Walker syndrome
- (B) Fanconi-Bickel syndrome
- (C) Beckwith syndrome
- (D) Menke's disease

• • •

**40.** Citrate used in fatty acid synthesis uses which enzyme?

- (A) Citrate synthase
- (B) ATP citrate lyase
- (C) Aconitase
- (D) Malic enzyme

• • •

**41.** Regarding apoprotein-C, which of the following is true?

- (A) Activates lipoprotein lipase
- (B) Inactivates lipoprotein lipase
- (C) Facilitates triglyceride transport
- (D) All of the above

• • •

**42.** What will you give in the diet to stop chyluria?

- (A) Short chain fatty acids
- (B) Medium chain fatty acids
- (C) Long chain fatty acids
- (D) Omega-3 unsaturated fatty acids

• • •

**43.** In Zellweger syndrome, there is accumulation of which type of fatty acids?

- (A) Accumulation of long fatty acids
- (B) Accumulation of short chain fatty acids
- (C) Accumulation of very long chain fatty acids
- (D) Accumulation of medium chain fatty acids

• • •

**44.** Selenocysteine is associated with which of the following?

- (A) Carbonic anhydrase
- (B) Catalase
- (C) Deiodinase
- (D) Transferase

• • •

**45.** The urea cycle and the Krebs cycle are linked at which intermediate?

- (A) Arginine
- (B) Ornithine
- (C) Oxaloacetate
- (D) Fumarate

• • •

**46.** Light of 250 nm wavelength is absorbed by which amino acid?

- (A) Arginine
- (B) Alanine
- (C) Tyrosine
- (D) Histidine

• • •

**47.** Why is the citric acid cycle called an amphibolic pathway?

- (A) Both exergonic and endergonic reactions take place
- (B) Metabolites are utilized in other pathways
- (C) It can proceed both in forward and backward direction
- (D) Same enzymes can be used in reverse directions

• • •

**48.** Which one of the following statements concerning gluconeogenesis is correct?

- (A) It occurs in muscle
- (B) It is stimulated by fructose 2,6-bisphosphate
- (C) It is inhibited by elevated levels of acetyl CoA
- (D) It is important in maintaining blood glucose during the normal overnight fast

• • •

**49.** During overnight fasting, what occurs?

- (A) Glucose decreases
- (B) FFA increases
- (C) Increased gluconeogenesis
- (D) Increased beta-hydroxybutyrate

• • •

**50.** The coenzyme form of pyridoxine (vitamin B6) is?

- (A) ADP
- (B) NAD
- (C) PLP
- (D) FAD

• • •

**51.** Uric acid is formed by?

- (A) Catabolism of proteins
- (B) Catabolism of ketones
- (C) Catabolism of purines
- (D) Catabolism of pyrimidines

• • •

**52.** A nucleotide consists of all of the following except?

- (A) Sugar
- (B) Phosphate
- (C) Fatty acid
- (D) Base

• • •

**53.** The function of a primer is in?

- (A) Transcription
- (B) Translation
- (C) Initiation of DNA replication
- (D) Termination of DNA replication

• • •

**54.** Which of the following usually requires an RNA intermediate for cloning or replication?

- (A) Transposons
- (B) Plasmids
- (C) Phages
- (D) Cosmids

• • •

**55.** The folds in collagen are due to?

- (A) Glycine
- (B) Alanine
- (C) Arginine
- (D) Histidine

• • •

**56.** This molecule attaches to a protein before its destruction?

- (A) Ubiquitin
- (B) RNase F
- (C) Zymase
- (D) Chaperone

• • •

**57.** Which isoform of LDH is raised in anemia?

- (A) LDH 5
- (B) LDH 4
- (C) LDH 3
- (D) LDH 2

• • •

**58.** Cell-matrix adhesions are mediated by?

- (A) Cadherins
- (B) Integrins
- (C) Selectins
- (D) Calmodulin

• • •

**59.** Regarding neurofibromatosis, all are true except:

- (A) Autosomal recessive
- (B) Associated with cataract
- (C) Scoliosis
- (D) Multiple fibroma

• • •

**60.** Regarding desmoid tumour, which is not correct?

- (A) Often seen below the umbilicus
- (B) Unencapsulated
- (C) More common in women
- (D) Highly radiosensitive

• • •

**61.** Most common cause of dissecting hematoma is because of:

- (A) Hypertension
- (B) Marfan's syndrome
- (C) Iatrogenic
- (D) Kawasaki disease

• • •

**62.** Creola bodies are seen in:

- (A) Bronchial asthma
- (B) Chronic bronchitis
- (C) Emphysema
- (D) Bronchiectasis

• • •

**63.** Renal papillary necrosis is almost always associated with one of the following conditions:

- (A) Diabetes mellitus
- (B) Analgesic nephropathy
- (C) Chronic pyelonephritis
- (D) Post streptococcal glomerulonephritis

• • •

**64.** Starry sky appearance is seen in -

- (A) Burkitt lymphoma
- (B) CLL
- (C) Diffuse large B cell lymphoma
- (D) ALCL

• • •

**65.** Russell bodies are found in -

- (A) Multiple myeloma
- (B) Gonadal tumor
- (C) Parkinsonism
- (D) Intracranial neoplasms

• • •

**66.** The anaemia associated with leukaemia is?

- (A) Iron deficiency
- (B) Megaloblastic type
- (C) Myelophthistic type
- (D) All of the above

• • •

**67.** Pathological manifestation of chronic alcoholism includes all of the following except -

- (A) Piecemeal necrosis
- (B) Ballooning degeneration
- (C) Microvesicular fatty changes
- (D) Central hyaline sclerosis

• • •

**68.** What is the histological appearance of brain in Creutzfeldt-Jakob disease -

- (A) Neuronophagia
- (B) Spongiform change in brain
- (C) Microabscesses
- (D) Demyelination

• • •

**69.** Most common malignancy of fallopian tube

- (A) Squamous cell carcinoma
- (B) Serous carcinoma
- (C) Teratoma
- (D) Choriocarcinoma

• • •

**70.** Glomus tumor is seen in -

- (A) Retroperitoneum
- (B) Soft tissue
- (C) Distal portion of digits
- (D) Proximal portion of digits

• • •

**71.** Type B adverse drug reaction is -

- (A) Augmented effect of drug
- (B) Unpredictable bizarre reaction
- (C) Effect seen on chronic use of drug
- (D) Delayed effect of drug

• • •

**72.** Which of the following causes increased renin on prolonged use?

- (A) Clonidine
- (B) Enalapril
- (C) Methyldopa
- (D) Beta blocker

• • •

**73.** Which drug is not included among DMARDs?

- (A) Chloroquine
- (B) Vincristine
- (C) Azathioprine
- (D) Leflunomide

• • •

**74.** Desmopressin is preferred over vasopressin because desmopressin -

- (A) More potent
- (B) More selective for V1 receptor
- (C) Has little vasoconstrictor activity
- (D) a and c

• • •

**75.** Which is a short acting non-depolarizing neuromuscular blocker?

- (A) Rocuronium
- (B) Suxamethonium
- (C) Mivacurium
- (D) Pancuronium

• • •

**76.** Finasteride is a:

- (A) 5 alpha reductase inhibitor
- (B) PDE inhibitor
- (C) Alpha 1a blocker
- (D) Androgen receptor blocker

• • •

**77.** An 86 year old lady presented with severe constipation. She was a known hypertensive on medications for 10 years. In clinic, her BP was 157/98 mm Hg with a heart rate of 58/min. On taking her BP in the supine position it was found to be 90/60 mm Hg. She had a recent history of depression. She is taking atenolol, thiazide, imipramine, haloperidol and docusate. What will be the next best step in the management?

- (A) Change atenolol and thiazide to a calcium channel blocker and ACE inhibitor and add bisacodyl for constipation
- (B) Change imipramine and haloperidol to fluoxetine and risperidone and add bisacodyl for constipation
- (C) Only add bisacodyl for constipation and continue rest of the medications
- (D) Discontinue all her medications and start her on steroids

• • •

**78.** Longest acting sulphonamide is -

- (A) Sulfadiazine
- (B) Sulphadoxine
- (C) Sulfamethoxazole
- (D) Sulfamethiazole

• • •

**79.** Sulphonamide injection causes decrease in folic acid by?

- (A) Competitive inhibition
- (B) Non competitive inhibition
- (C) Uncompetitive inhibition
- (D) Allosteric inhibition

• • •

**80.** All are true regarding Sunitinib except -

- (A) It inhibits tyrosine kinase receptors
- (B) It is used for renal cell carcinoma
- (C) It is used for the treatment of GIST
- (D) It is excreted primarily in urine

• • •

**81.** Plaintiff is a person who?

- (A) Files case in civil court
- (B) Acts as defender
- (C) Gives judgement
- (D) None

• • •

**82.** Which of the following is not seen in finger prints?

- (A) Loop
- (B) Circles
- (C) Whorl
- (D) Arch

• • •

**83.** In medicolegal autopsy, cavity to be opened first is?

- (A) Thoracic
- (B) Abdomen
- (C) Cervical
- (D) Any of the above

• • •

**84.** Marshall's triad is seen in?

- (A) Explosive injury
- (B) Gunshot injury
- (C) Drowning injury
- (D) None

• • •

**85.** A victim was choked with the mouth and nose covered, while the neck was held in the bend of the assailant's elbow from behind. This method of strangulation is called?

- (A) Mugging
- (B) Garroting
- (C) Bansdola
- (D) None of the above

• • •

**86.** Which of the following substances acts as both a poison and an antidote?

- (A) Copper sulphate ( $\text{CuSO}_4$ )
- (B) Mercuric chloride ( $\text{HgCl}_2$ )
- (C) Arsenic trioxide ( $\text{As}_2\text{O}_3$ )
- (D) Thallium arsenate

• • •

**87.** Muttering delirium is characteristically seen in poisoning with:

- (A) Ricinus (castor)
- (B) Dhatura (Datura)
- (C) Cocaine
- (D) Aconite

• • •

**88.** The term vitriolage means:

- (A) Using vitriol for suicide
- (B) Using vitriol for murder
- (C) Vitriol throwing
- (D) Perforation of the stomach caused by  $\text{H}_2\text{SO}_4$

• • •

**89.** Who is regarded as the Father of Toxicology?

- (A) Paracelsus
- (B) Galen
- (C) Galton
- (D) Orfila

• • •

**90.** Lendrum stain (Phloxine-Tartrazine) is done at autopsy for the diagnosis of?

- (A) Air embolism
- (B) Fat embolism
- (C) Amniotic fluid embolism
- (D) None

• • •

**91.** In nutrient agar, the concentration of agar is:

- (A) 1%
- (B) 1.5%
- (C) 3%
- (D) 4%

• • •

**92.** A child comes with fever, cold, cough and a membrane over the tonsils. A nasal swab is taken. Culture should be done on which medium for the earliest diagnosis?

- (A) Loeffler's serum slope
- (B) L. J. medium
- (C) MacConkey's agar
- (D) Citrate medium

• • •

**93.** Cholera toxin is due to -

- (A) Chromosome
- (B) Plasmid
- (C) Phage
- (D) Transposons

• • •

**94.** A child with fever, red blood cells (RBCs) and pus in the stool. The causative organism is?

- (A) ETEC (Enterotoxigenic E. coli)
- (B) EHEC (Enterohaemorrhagic E. coli)
- (C) EPEC (Enteropathogenic E. coli)
- (D) EAEC (Enteroaggregative E. coli)

• • •

**95.** Reactive tubercular arthritis is known as:

- (A) Spina ventosa
- (B) Pott's disease
- (C) Poncet's disease
- (D) None

• • •

**96.** In malaria, the sexual cycle is -

- (A) Sporozoite to gametocytes
- (B) Gametocytes to sporozoite
- (C) Occurs in humans
- (D) Responsible for relapse

• • •

**97.** Toxoplasma in children causes:

- (A) Chorioretinitis
- (B) Conjunctivitis
- (C) Keratitis
- (D) Papillitis

• • •

**98.** Which of the following is a primary cell line?

- (A) Chick embryo fibroblast
- (B) HeLa cells
- (C) Vero cells
- (D) WI-38

• • •

**99.** Bollinger bodies are seen in?

- (A) Chickenpox
- (B) Cowpox
- (C) Fowlpox
- (D) Smallpox

• • •

**100.** Binding of gp120 causes:

- (A) Infection of target cell
- (B) Facilitation of co-receptor
- (C) Fusing of virus and target cell
- (D) None

• • •

**101.** Prozone phenomenon is seen with?

- (A) Same concentration of antibody and antigen
- (B) Antigen excess to antibody
- (C) Antibody excess to antigen
- (D) Hyperimmune reaction

• • •

**102.** Which of the following T cell independent antigen acts through?

- (A) T-cell
- (B) B-cell
- (C) Macrophages
- (D) CD8+ T cells

• • •

**103.** Complement components formed in the liver are?

- (A) C2, C4
- (B) C3, C6, C9
- (C) C5, C8
- (D) C1

• • •

**104.** An 18 year old girl presents with watery diarrhea. The most likely causative agent is?

- (A) Rotavirus
- (B) V. cholerae
- (C) Salmonella
- (D) Shigella

• • •

**105.** Use of iodized salt in the iodine deficiency control programme is an example of?

- (A) Primary prevention
- (B) Secondary prevention
- (C) Tertiary prevention
- (D) None

• • •

**106.** Which study is best to establish the definitive cause of a disease?

- (A) Case-control
- (B) Cohort
- (C) Ecological
- (D) Cross-sectional

• • •

**107.** The time interval between receipt of infection and the point of maximum infectivity of the host is known as?

- (A) Incubation period
- (B) Serial interval
- (C) Generation time
- (D) Communicable period

• • •

**108.** Mass vaccination is ineffective in which of the following?

- (A) Measles
- (B) Polio
- (C) Tetanus
- (D) None

• • •

**109.** At what age is the first dose of measles vaccination given?

- (A) 9 months
- (B) At birth
- (C) 4 weeks
- (D) 8 weeks

• • •

**110.** What is the amount of diphtheria toxoid present in the pediatric DT vaccine?

- (A) 5 Lf
- (B) 10 Lf
- (C) 15 Lf
- (D) 25 Lf

• • •

**111.** What is the osmolarity of the new (reduced osmolarity) ORS in mmol per litre?

- (A) 270
- (B) 245
- (C) 290
- (D) 310

• • •

**112.** In leprosy, a duration of 2 years is relevant with regard to which of the following?

- (A) Treatment of paucibacillary leprosy
- (B) Treatment of multibacillary leprosy
- (C) Post-treatment surveillance of paucibacillary leprosy
- (D) Post-treatment surveillance of multibacillary leprosy

• • •

**113.** Which of the following is the most common cancer in men?

- (A) Bladder cancer
- (B) Colorectal cancer
- (C) Prostate cancer
- (D) Oral cancer

• • •

**114.** Net reproduction rate is defined as which of the following?

- (A) Number of live births per 1000 mid-year population
- (B) Number of live births per 1000 women of child bearing age
- (C) Number of daughters a newborn girl will have during her life time
- (D) None of the above

• • •

**115.** Which of the following is the least common cause of infant mortality in India?

- (A) Infections
- (B) Prematurity
- (C) Birth injuries
- (D) Congenital malformations

• • •

**116.** What is the approximate total daily energy requirement in late pregnancy?

- (A) 2500 cal
- (B) 1400 cal
- (C) 1000 cal
- (D) 1800 cal

• • •

**117.** Which toxin is responsible for epidemic dropsy?

- (A) BOAA
- (B) Aflatoxin
- (C) Sanguinarine
- (D) Pyrrolizidine

• • •

**118.** Which is the recommended measure for the iodine deficiency control programme?

- (A) Health education
- (B) Water testing
- (C) Fortification of salt
- (D) None

• • •

**119.** What amount of bleaching powder is necessary to disinfect choleric stools?

- (A) 50 gm/lit
- (B) 75 gm/lit
- (C) 90 gm/lit
- (D) 100 gm/lit

• • •

**120.** Which is the most common heavy metal poisoning in the world?

- (A) Lead
- (B) Arsenic
- (C) Mercury
- (D) Cadmium

• • •

**121.** What is the approximate amount of infectious waste produced in hospitals (as a fraction of total hospital waste)?

- (A) 45%
- (B) 65%
- (C) 80%
- (D) 100%

• • •

**122.** In a left (negatively) skewed distribution curve, which statement is true?

- (A) Mean = Median
- (B) Mean < Mode
- (C) Mean > Mode
- (D) Mean = Mode

• • •

**123.** The Ujjwala scheme is meant for the prevention of which problem?

- (A) Child abuse
- (B) Child trafficking
- (C) Child labour
- (D) None

• • •

**124.** Above which level of the heat stress index (HSI) is it not possible to work comfortably?

- (A) 20 - 40
- (B) 40 - 60
- (C) 60 - 80
- (D) 80 - 100

• • •

**125.** The visual axis of the eye is best defined as the line from the:

- (A) Center of cornea to retina
- (B) Object to fovea
- (C) Center of lens to cornea
- (D) None

• • •

**126.** Which of the following are features of vernal keratoconjunctivitis?

- (A) Shield ulcer
- (B) Horner-Trantas spots
- (C) Papillary hypertrophy
- (D) All of the above

• • •

**127.** Disciform keratitis is seen in infection with which of the following?

- (A) HSV
- (B) HIV
- (C) HBV
- (D) Rubella

• • •

**128.** What is the most common cause of anterior uveitis?

- (A) CMV
- (B) Ankylosing spondylitis
- (C) Toxoplasma
- (D) None

• • •

**129.** Second sight is seen in which type of cataract?

- (A) Nuclear cataract
- (B) Cortical cataract
- (C) Zonular cataract
- (D) Punctate cataract

• • •

**130.** Laser iridotomy is done in which type of glaucoma?

- (A) Angle closure glaucoma
- (B) Open angle glaucoma
- (C) Pigmentary glaucoma
- (D) None

• • •

**131.** What is the characteristic deposit seen in retinal macular degeneration?

- (A) Iron
- (B) Drusen
- (C) Lipochrome
- (D) Hemosiderin

• • •

**132.** All of the following are seen in third (3rd) cranial nerve palsy EXCEPT?

- (A) Mydriasis
- (B) Loss of light reflex
- (C) Loss of abduction
- (D) Ptosis

• • •

**133.** Which of the following is the longest extraocular muscle?

- (A) Superior rectus (SR)
- (B) Medial rectus (MR)
- (C) Superior oblique (SO)
- (D) Inferior oblique (IO)

• • •

**134.** The most common orbital tumor has its origin from which structure?

- (A) Blood vessels
- (B) Nerves
- (C) Muscle
- (D) Lymph node

• • •

**135.** An abnormally eccentric placed pupil is called?

- (A) Polycoria
- (B) Corectopia
- (C) Aniridia
- (D) Ectopia lentis

• • •

**136.** Iris coloboma is most commonly located in which quadrant?

- (A) Inferotemporal
- (B) Superotemporal
- (C) Inferonasal
- (D) Superonasal

• • •

**137.** A left axis deviation of  $-30^\circ$  to  $-60^\circ$  on the ECG is seen in?

- (A) Left ventricular hypertrophy
- (B) Right ventricular hypertrophy
- (C) Aortic stenosis
- (D) Left atrial enlargement

• • •

**138.** A person has asthma attacks more than once during the day and at least once during the night. This is classified as?

- (A) Mild intermittent asthma
- (B) Mild persistent asthma
- (C) Moderate asthma
- (D) Severe asthma

• • •

**139.** The most common presentation of IgA nephropathy is?

- (A) Nephrotic syndrome
- (B) Nephritic syndrome
- (C) Microscopic hematuria
- (D) Repeated gross hematuria

• • •

**140.** All of the following are used for treatment of H. pylori, except?

- (A) Gentamycin
- (B) Clarithromycin
- (C) Metronidazole
- (D) Amoxicillin

• • •

**141.** Secretory diarrhea is NOT seen in which of the following conditions?

- (A) Phenolphthalein
- (B) Celiac disease
- (C) Cholera
- (D) Addison's disease

• • •

**142.** Predominantly sensory neuropathy is/are caused by:

- (A) Cisplatin
- (B) Pyridoxine excess
- (C) Suramin
- (D) a and b

• • •

**143.** Regarding herpes encephalitis, all are true EXCEPT:

- (A) Focal symptoms common
- (B) Temporal lobe involved
- (C) MRI is diagnostic
- (D) EEG not diagnostic

• • •

**144.** The immunoglobulin most commonly involved in multiple myeloma is:

- (A) IgG
- (B) IgM
- (C) IgA
- (D) IgD

• • •

**145.** The most classical symptom of VIPoma is:

- (A) Gall stones
- (B) Secretory diarrhea
- (C) Steatorrhea
- (D) Flushing

• • •

**146.** Antibodies most commonly seen in drug-induced lupus are:

- (A) Anti-ds DNA antibodies
- (B) Anti-Sm antibodies
- (C) Anti-Ro antibodies
- (D) Antihistone antibodies

• • •

**147.** Muir-Torre syndrome shows:

- (A) Sebaceous keratomas
- (B) Lisch nodules
- (C) Intestinal polyp
- (D) Hyperelastic joints

• • •

**148.** All of the following are risk factors for deep vein thrombosis (DVT) except -

- (A) Duration of surgery more than thirty minutes
- (B) Obesity
- (C) Age less than forty years
- (D) Use of the oestrogen-progesterone contraceptive pills

• • •

**149.** Inverted Champagne bottle appearance is seen in -

- (A) Varicose veins
- (B) DVT
- (C) Lipodermatosclerosis
- (D) Venous ulceration

• • •

**150.** Which of the following findings in ascitic fluid is diagnostic of spontaneous bacterial peritonitis?

- (A) Neutrophils  $> 250/\text{mm}^3$
- (B) WBC  $< 100/\text{mL}$
- (C) Ascitic lactate level  $< 25 \text{ mg/dL}$
- (D) Ascitic fluid pH  $> 7.35$

• • •

**151.** Ogilvie's syndrome most commonly involves -

- (A) Stomach
- (B) Colon
- (C) Gallbladder
- (D) Small intestine

• • •

**152.** Raspberry tumor is seen in -

- (A) Umbilical fistula
- (B) Meckel's diverticulum
- (C) Umbilical adenoma
- (D) Umbilical granuloma

• • •

**153.** In injury to the penis, which of the following prevents extravasation of blood -

- (A) Buck's fascia
- (B) Fascia of Camper
- (C) Fascia transversalis
- (D) None

• • •

**154.** Which of the following is FALSE about hepatic adenoma?

- (A) Benign lesion
- (B) OCP use
- (C) Older females
- (D) Cold on isotope scan

• • •

**155.** VIPoma is associated with which syndrome?

- (A) Zollinger Ellison
- (B) Verner Morrison
- (C) Carcinoid syndrome
- (D) Cushing's syndrome

• • •

**156.** In Marfan's syndrome, aortic aneurysm occurs most commonly in:

- (A) Ascending aorta
- (B) Descending aorta
- (C) Abdominal aorta
- (D) Arch of aorta

• • •

**157.** Dosage of intravenous fluid for a 2 month old child with diarrhoea and severe dehydration:

- (A) 100 ml/kg in 6 hours
- (B) 50 ml/kg in 6 hours
- (C) 75 ml/kg in 6 hours
- (D) 80 ml/kg in 6 hours

• • •

**158.** LEOPARD syndrome includes all except:

- (A) Growth retardation
- (B) ECG changes
- (C) Hypertelorism
- (D) Hypergonadism

• • •

**159.** Most common cerebellar tumour in children?

- (A) Astrocytoma
- (B) Medulloblastoma
- (C) Ependymoma
- (D) DNET

• • •

**160.** Upper segment to lower segment ratio in a 3 year old child is:

- (A) 1.2
- (B) 1.3
- (C) 1.4
- (D) 1.6

• • •

**161.** Diarrhoea in a child of 12 months, the dose of zinc is?

- (A) 1 mg / 10 to 14 days
- (B) 10 mg / 10 to 14 days
- (C) 15 mg / 10 to 14 days
- (D) 20 mg / 10 to 14 days

• • •

**162.** A baby is born at 27 weeks of gestation, required mechanical ventilation for the next 4 weeks and oxygen for the next 1 week. He was maintained at room air subsequently. As per the new Bronchopulmonary dysplasia (BPD) definition, he has which of the following?

- (A) Mild BPD
- (B) Moderate BPD
- (C) Severe BPD
- (D) No BPD

• • •

**163.** Which of the following statements regarding fetal circulation is correct?

- (A) The liver and heart of the fetus receive blood with very high oxygen saturation
- (B) PO<sub>2</sub> of fetal blood leaving the placenta is slightly greater than maternal mixed venous PO<sub>2</sub>
- (C) The presence of fetal hemoglobin shifts the oxyhemoglobin dissociation to the right
- (D) The foramen ovale closes during the third trimester unless the fetus has an atrial septal defect

• • •

**164.** Cause of conjugated hyperbilirubinemia is?

- (A) Rotor syndrome
- (B) Breast milk jaundice
- (C) Crigler-Najjar syndrome
- (D) Gilbert syndrome

• • •

**165.** Highest cure rate is of:

- (A) Wilms tumor
- (B) Retinoblastoma
- (C) Rhabdomyosarcoma
- (D) All

• • •

**166.** The MOST common type of dislocation of the elbow joint is:

- (A) Posterior
- (B) Posterolateral
- (C) Posteromedial
- (D) Lateral

• • •

**167.** Most common complication of fracture neck of femur is:

- (A) Malunion
- (B) Avascular necrosis (AVN)
- (C) Nonunion
- (D) Arthritis

• • •

**168.** Partial anterior dislocation of one segment of the spine over another is:

- (A) Spondylosis
- (B) Spondylolisthesis
- (C) Kyphosis
- (D) Scoliosis

• • •

**169.** In a child with suspected developmental dysplasia of the hip (DDH), what is the investigation of choice?

- (A) X-ray
- (B) MRI
- (C) USG
- (D) CT scan

• • •

**170.** Which is the most common organism causing infection after an open fracture?

- (A) Pseudomonas
- (B) Staphylococcus aureus
- (C) Klebsiella
- (D) Gonococcus

• • •

**171.** In tuberculosis of the spine, which is the most common site of involvement?

- (A) Sacral
- (B) Cervical
- (C) Dorsolumbar
- (D) Lumbosacral

• • •

**172.** In Volkmann's ischaemic contracture, which artery is involved?

- (A) Radial
- (B) Brachial
- (C) Ulnar
- (D) Interosseous

• • •

**173.** Why should a fracture of the shaft of the femur be stabilised early?

- (A) To prevent blood loss
- (B) ARDS
- (C) Non union
- (D) Compartment syndrome

• • •

**174.** Hydrops fetalis is due to?

- (A) Rh mismatch
- (B) Hyperproteinemia
- (C) Placental hypoplasia
- (D) All of the above

• • •

**175.** What is the most common type of degeneration seen in uterine fibroids (leiomyomas)?

- (A) Calcareous
- (B) Hyaline
- (C) Red
- (D) Cystic

• • •

**176.** Which of the following is true regarding fallopian tube tuberculosis?

- (A) Most common type of genital TB
- (B) Size of the tubes is unchanged
- (C) Is asymptomatic
- (D) Primary focus of infection is always in the fallopian tubes

• • •

**177.** Regarding intrahepatic cholestasis of pregnancy, the FALSE statement is?

- (A) Bilirubin level  $> 2$  mg%
- (B) Most common cause of jaundice in pregnancy
- (C) Oestrogen is involved
- (D) Manifestations usually appear in the last trimester

• • •

**178.** Which of the following pelvic measurements is most commonly used clinically?

- (A) Diagonal conjugate
- (B) Anteroposterior diameter of the inlet
- (C) Transverse diameter of the outlet
- (D) Oblique diameter of the pelvis

• • •

**179.** Maximum chance of transmission to the baby during delivery is seen with which infection?

- (A) HSV (Herpes simplex virus)
- (B) CMV (Cytomegalovirus)
- (C) VZV (Varicella zoster virus)
- (D) Rubella

• • •

**180.** What is the preferred treatment for menorrhagia in the reproductive age group?

- (A) NOVA T (copper IUCD)
- (B) Cu IUD (copper intrauterine device)
- (C) OCPs (oral contraceptive pills)
- (D) Hysterectomy

• • •

**181.** A 45 year old female patient underwent hysterectomy. On the 7th post-operative day she complains of continuous dribbling of urine and fever. Micturition was not voluntary. What is the diagnosis?

- (A) Vesicovaginal fistula
- (B) Ureterovaginal fistula
- (C) Vesicouterine fistula
- (D) Urethrovaginal fistula

• • •

**182.** Regarding Hegar's sign, all are true EXCEPT?

- (A) It is elicited by the bimanual palpation method
- (B) It is difficult to elicit in obese women
- (C) It can be done at 14 weeks
- (D) It is present in about two-thirds of cases

• • •

**183.** In a non-stress test (NST), all of the following are assessed except?

- (A) Variability
- (B) Acceleration
- (C) Time period
- (D) Oxytocin

• • •

**184.** During pregnancy, cardiac output increases to its maximum at which week?

- (A) 26-28 weeks
- (B) 30-32 weeks
- (C) 32-34 weeks
- (D) 34-36 weeks

• • •

**185.** What is the presenting part in a transverse lie?

- (A) Shoulder
- (B) Face
- (C) Vertex
- (D) Brow

• • •

**186.** Which of the following is NOT a nevus of melanocytic origin?

- (A) Mongolian spot
- (B) Nevus of Ito
- (C) Nevus of Ota
- (D) Becker nevus

• • •

**187.** Oculo-oro-genital ulcers are a feature of which condition?

- (A) Behcet disease
- (B) Lichen planus
- (C) SLE
- (D) Psoriasis

• • •

**188.** The levels (stages) of ether anaesthesia were demonstrated by whom?

- (A) Morton
- (B) Guedel
- (C) Thompson
- (D) None

• • •

**189.** Which anaesthetic agent is neither metabolised by the liver nor by the kidney?

- (A) Atracurium
- (B) Vecuronium
- (C) Pancuronium
- (D) Rocuronium

• • •

**190.** Which is the inducing agent of choice in a patient presenting with shock?

- (A) Isoflurane
- (B) Desflurane
- (C) Ketamine
- (D) Thiopentone

• • •

**191.** Which drug is used for analgesia in OPD or day-care (outpatient) surgical settings?

- (A) Morphine
- (B) Pethidine
- (C) Fentanyl
- (D) Alfentanil

• • •

**192.** The Hounsfield unit (HU) is a measure used in which imaging modality?

- (A) CT
- (B) MRI
- (C) PET
- (D) USG

• • •

**193.** Corkscrew esophagus on a barium study is characteristically seen in which of the following conditions?

- (A) Carcinoma esophagus
- (B) Scleroderma
- (C) Achalasia cardia
- (D) Diffuse esophageal spasm

• • •

**194.** What is the investigation of choice for screening of proximal internal carotid artery stenosis?

- (A) Doppler flow USG
- (B) CT subtraction angiography
- (C) MRI
- (D) Angiography (DSA)

• • •

**195.** What is the physical half-life of Iodine-131?

- (A) 8 hours
- (B) 8 days
- (C) 8 weeks
- (D) 8 months

• • •

**196.** Formication (a sensation of insects crawling on or under the skin) is characteristically seen with which of the following?

- (A) Acute amphetamine intoxication
- (B) Chronic use of amphetamine
- (C) Alcohol withdrawal
- (D) Cannabis poisoning

• • •

**197.** In depression, there is a deficiency of which neurotransmitter?

- (A) 5-HT
- (B) Acetylcholine
- (C) Dopamine
- (D) GABA

• • •

**198.** Persistent preoccupation with having a serious illness despite normal body function is called?

- (A) Obsession
- (B) Somatization
- (C) Hypochondriasis
- (D) Conversion disorder

• • •

**199.** How do you differentiate between psychological and organic erectile dysfunction?

- (A) Nocturnal penile tumescence
- (B) PIPE therapy
- (C) Sildenafil induced erection
- (D) Squeeze technique

• • •

200.



The type of connective tissue present in the arrow-marked area is:

- (A) Loose and irregular
- (B) Specialized
- (C) Dense irregular
- (D) Dense regular